## References

- K. T. Fang, D. K. J. Lin, P. Winker and Y. Zhang (2000), Uniform Design: Theory and Application, Technometrics, 42, 237–248.
- [2] K. T. Fang, X. Lu and P. Winker (2003), Lower bounds for centered and wrap-around  $L_2$ -discrepancies and construction of uniform design by threshold accepting, Journal of Complexity, **19**, 692–711.
- [3] K. T. Fang, C. X. Ma and P. Winker (2002), Centered L<sub>2</sub>-discrepancy of random sampling and Latin hypercube design, and construction of uniform designs, Mathematics of Computation, 71, 275–296.
- [4] K. T. Fang, Y. Tang and J. X. Yin (2004), Lower bounds for wrap-around L<sub>2</sub>-discrepancy and constructions of symmetrical uniform designs, Technical Report MATH-372, Hong Kong Baptist University.
- [5] K. T. Fang, D. Maringer, Y. Tang and P. Winker (2004), Lower Bounds and stochastic optimization algorithms for uniform design with three and four level, working paper.
- [6] F. J. Hickernell (1998a), A generalized discrepancy and quadrature error bound, Mathematics of Computation, 67, 299–322.
- [7] F. J. Hickernell (1998b), Lattice rules: how well do they measure up? in P. Hellekalek and G. Larcher (eds), Random and Quasi-Random Point Sets, Springer-Verlag, 106–166.
- [8] F. J. Hickernell and M. Q. Liu (2002), Uniform designs limit aliasing, Biometrika, 89, 893–904.
- [9] K. T. Fang, G. N. Ge and M. Q. Liu (2004), Construction of optimal supersaturated designs by the packing method, Science of China Ser. (A) 47, 128–143.
- [10] K. T. Fang, G. N. Ge, M. Q. Liu and H. Qin (2004), Combinatorial constructions for optimal supersaturated designs, Discrete Mathematics 279, 191–202.

- [11] K. T. Fang, X. Lu, Y. Tang and J. X. Yin (2004), Constructions of uniform designs by using resolvable packings and coverings, Discrete Mathematics, 274, 25–40.
- [12] K. T. Fang, Y. Tang and J. X. Yin, Resolvable partially pairwise balanced designs and their applications in computer experiments, to appear in Utilitas Mathematica.