



## Comment on “A Tuning-Free Robust and Efficient Approach to High-Dimensional Regression”

Jianqing Fan , Cong Ma & Kaizheng Wang

To cite this article: Jianqing Fan , Cong Ma & Kaizheng Wang (2020) Comment on “A Tuning-Free Robust and Efficient Approach to High-Dimensional Regression”, Journal of the American Statistical Association, 115:532, 1720-1725, DOI: [10.1080/01621459.2020.1837138](https://doi.org/10.1080/01621459.2020.1837138)

To link to this article: <https://doi.org/10.1080/01621459.2020.1837138>



Published online: 18 Dec 2020.



Submit your article to this journal [↗](#)



Article views: 52



View related articles [↗](#)



View Crossmark data [↗](#)



# Comment on “A Tuning-Free Robust and Efficient Approach to High-Dimensional Regression”

Jianqing Fan<sup>a</sup>, Cong Ma<sup>b</sup>, and Kaizheng Wang<sup>c</sup>

<sup>a</sup>Department of Operations Research and Financial Engineering, Princeton University, Princeton, NJ; <sup>b</sup>Department of Electrical Engineering and Computer Sciences, UC Berkeley, Berkeley, CA; <sup>c</sup>Department of Industrial Engineering and Operations Research, Columbia University, New York, NY

We congratulate the authors for their important and timely contributions to robust and tuning-free high-dimensional linear regression. The proposed loss function originates from non-parametric rank-based estimation and enjoys certain pivotal properties that facilitate the selection of the tuning parameter. While tuning-free probably sounds over claimed, the proposed penalization parameter is indeed more interpretable, easier to select, and is independent of noise variance. We welcome the opportunity to make a few comments from various perspectives and discuss open questions that are worth studying.

## 1. Historical Perspectives on Rank Lasso

Consider the linear model

$$y = X\beta_0 + \varepsilon, \quad (1)$$

where  $y = (y_1, \dots, y_n)^\top \in \mathbb{R}^n$  is a response vector,  $X = (x_1, \dots, x_n)^\top \in \mathbb{R}^{n \times p}$  is a design matrix,  $\beta_0 \in \mathbb{R}^p$  is an unknown coefficient vector, and  $\varepsilon = (\varepsilon_1, \dots, \varepsilon_n)^\top \in \mathbb{R}^n$  is a noise vector with iid entries. The Rank Lasso estimator (Wang, Peng, et al. 2020) for estimating  $\beta_0$  is given by

$$\hat{\beta}(\lambda) = \operatorname{argmin}_{\beta \in \mathbb{R}^p} \{L_n(\beta) + \lambda \|\beta\|_1\}, \quad (2)$$

where the empirical loss function  $L_n(\beta)$  is defined as

$$L_n(\beta) = \frac{1}{n(n-1)} \sum_{i \neq j} |(y_i - x_i^\top \beta) - (y_j - x_j^\top \beta)|, \quad (3)$$

and  $\lambda \geq 0$  is a regularization parameter to be chosen. This is equivalent to fitting the linear model

$$y_i - y_j = (x_i - x_j)^\top \beta_0 + (\varepsilon_i - \varepsilon_j) \quad (4)$$

using  $\ell_1$  regression, since the error distribution  $\varepsilon_i - \varepsilon_j$  is symmetric.

The proposed Rank Lasso estimator (2) is an extension of the traditional nonparametric rank-based estimator for linear models to the high-dimensional setting (see Hettmansperger and McKean 1978, 2010). Without the  $\ell_1$  penalty, a general rank-based estimator takes the following form

$$\hat{\beta} = \operatorname{argmin}_{\beta \in \mathbb{R}^p} \left\{ \sum_{i=1}^n a \left( R \left( y_i - x_i^\top \beta \right) \right) \left( y_i - x_i^\top \beta \right) \right\}, \quad (5)$$

where  $R(y_i - x_i^\top \beta)$  denotes the rank of  $y_i - x_i^\top \beta$  among the residuals  $\{y_i - x_i^\top \beta\}_{1 \leq i \leq n}$  and  $a(i) = \phi(i/(n+1))$  for some nondecreasing function  $\phi(\cdot) : (0, 1) \mapsto \mathbb{R}$  satisfying  $\int \phi(u) du = 0$  and  $\int \phi^2(u) du = 1$ . An illustrative example of  $\phi(\cdot)$  is the sign function, that is,  $\phi(u) = \operatorname{sgn}(u - 1/2)$ . Under this circumstance, the general rank-based estimator (5) recovers the well-known least absolute deviation estimator.

In fact, Rank Lasso corresponds to the rank-based estimator with the Wilcoxon score function  $\phi(u) = \sqrt{12}(u - 1/2)$ , that is,

$$L_n(\beta) = \frac{2(n+1)}{\sqrt{3n(n-1)}} \sum_{i=1}^n a \left( R \left( y_i - x_i^\top \beta \right) \right) \left( y_i - x_i^\top \beta \right).$$

For future convenience, we denote  $D(y - X\beta) = \sum_{i=1}^n a(R(y_i - x_i^\top \beta))(y_i - x_i^\top \beta)$  and rewrite the Rank Lasso estimator as

$$\hat{\beta} = \operatorname{argmin}_{\beta \in \mathbb{R}^p} \left\{ \frac{2(n+1)}{\sqrt{3n(n-1)}} D(y - X\beta) + \lambda \|\beta\|_1 \right\}. \quad (6)$$

## 2. Computational Efficiency and Effect of Subsampling

Denote by  $f(\cdot)$  the density of the error distribution. The relative efficiency between the rank regression and the  $\ell_1$ -regression for symmetric error distributions is  $3[\int f^2(u) du]^2 / f(0)^2$  (Hodges and Lehmann 1963), which is 0.75, 1.04, 1.25, 1.37, 1.43, 1.50 for  $t_\nu$  distribution with  $\nu = 1, 2, 4, 8, 16, \infty$  (normal) and 3 for the uniform distribution. Namely, for a variety of distributions, the rank regression estimator has high efficiency in comparison with  $\ell_1$ -regression. Why, then, the  $\ell_1$ -regression is far more popular? One possible cause, as pointed out by the authors in Wang, Peng, et al. (2020, sec. 4.2), is the computational burden for obtaining the Rank Lasso estimator—the  $U$ -statistics structure in (3), which involves  $n(n-1)$  pairs of samples. To alleviate this issue, the authors suggest a subsampling mechanism to reduce the computational cost of the Rank Lasso estimator. More specifically,  $N = m \cdot n$  terms are sampled with replacement from  $n(n-1)/2$  terms in the loss function (3), where  $m$  can vary from 1 to  $(n-1)/2$ . A smaller  $m$  brings more computational benefits, but larger bias as well as variance. In other words, it loses statistical efficiency when compared with

using the full sample. In this section, we complement the article by investigating empirically the effect of the sampling budget  $m$ . In addition, we propose another principled subsampling mechanism for Rank Lasso that demonstrates superiority over the above mentioned one.

To motivate the new subsampling mechanism, we recall from Section 1 that the loss function  $L_n$  is equal to the  $\ell_1$  loss (absolute deviation) over symmetrized samples  $\{(\mathbf{x}_i - \mathbf{x}_j, y_i - y_j)\}_{i \neq j}$ . When  $n$  is even and  $\tau$  is a random permutation over  $[n]$ , a simplified loss with the same expectation is

$$\tilde{L}_n(\boldsymbol{\beta}; \tau) = \frac{2}{n} \sum_{i=1}^{n/2} |(y_{\tau(2i)} - \mathbf{x}_{\tau(2i)}^\top \boldsymbol{\beta}) + (y_{\tau(2i-1)} - \mathbf{x}_{\tau(2i-1)}^\top \boldsymbol{\beta})|, \tag{7}$$

which only concerns  $n/2$  iid samples  $\{(\mathbf{x}_{\tau(2i)} - \mathbf{x}_{\tau(2i-1)}, y_{\tau(2i)} - y_{\tau(2i-1)})\}_{i=1}^{n/2}$ . Minimizing  $\tilde{L}_n(\boldsymbol{\beta}; \tau) + \lambda \|\boldsymbol{\beta}\|_1$  amounts to  $\ell_1$ -penalized least absolute deviations, which can be studied using general results for sparse quantile regression by Belloni and Chernozhukov (2011) and Fan, Fan, and Barut (2014). Similar to the current article, they also propose to compute a pivotal quantity as the tuning parameter by simulation.

The loss in (7) based on nonoverlapping pairwise differences greatly facilitates computation while enjoying the same rate of convergence. Yet it may suffer from loss of efficiency. Consider the classical asymptotics with fixed  $p$ , diverging  $n$ , and random designs. Ignore the  $\ell_1$  penalty for a moment. Suppose that  $\varepsilon_i$  has density  $f$ , and  $\mathbb{E}(\mathbf{x}_i \mathbf{x}_i^\top) = \Sigma$ . Define  $\tilde{\varepsilon}_i = \varepsilon_{\tau(2i)} - \varepsilon_{\tau(2i-1)}$  and  $\tilde{\mathbf{x}}_i = \mathbf{x}_{\tau(2i)} - \mathbf{x}_{\tau(2i-1)}$ . Then the density function of  $\tilde{\varepsilon}_i$  is  $\tilde{f}(x) = (f * f)(x)$ , where  $*$  denotes convolution, and  $\mathbb{E}(\tilde{\mathbf{x}}_i \tilde{\mathbf{x}}_i^\top) = 2\Sigma$ . According to Bassett and Koenker (1978), the least absolute deviations estimator  $\tilde{\boldsymbol{\beta}}_n = \operatorname{argmin}_{\boldsymbol{\beta} \in \mathbb{R}^p} \tilde{L}_n(\boldsymbol{\beta}; \tau)$  satisfies

$$\begin{aligned} \sqrt{\frac{n}{2}}(\tilde{\boldsymbol{\beta}}_n - \boldsymbol{\beta}_0) &\xrightarrow{d} \mathcal{N}\left(\mathbf{0}, \frac{1}{[2\tilde{f}(0)]^2} \tilde{\Sigma}^{-1}\right) \\ &= \mathcal{N}\left(\mathbf{0}, \frac{1}{8[\int_{\mathbb{R}} f^2(x) dx]^2} \Sigma^{-1}\right). \end{aligned}$$

The asymptotic relative efficiency of  $\tilde{\boldsymbol{\beta}}_n$  with respect to the least squares estimate is  $4\sigma^2[\int_{\mathbb{R}} f^2(x) dx]^2$ , which is 1/3 of that for the estimator based on Jaeckel’s Wilcoxon-type dispersion function in Wang, Peng, et al. (2020). A natural idea to bridge this gap is to average the loss (7) first among a few permutations and then optimize the object. More specifically, for some positive integer  $m$ , one randomly draws  $2m$  permutations  $\{\tau_1, \tau_2, \dots, \tau_{2m}\}$  over  $[n]$  (so that we have  $mn$  pairs), and then seeks

$$\hat{\boldsymbol{\beta}} = \operatorname{argmin}_{\boldsymbol{\beta}} \left\{ \frac{1}{2m} \sum_{i=1}^{2m} \tilde{L}_n(\boldsymbol{\beta}; \tau_i) + \lambda \|\boldsymbol{\beta}\|_1 \right\}.$$

It turns out that this average nonoverlapping-pair estimator (ANOPE) bears deep connections to the Rank Lasso estimator. Indeed, the original symmetrized loss function  $L_n(\boldsymbol{\beta})$  satisfies

$$L_n(\boldsymbol{\beta}) = \frac{1}{n!} \sum_{\tau} \tilde{L}_n(\boldsymbol{\beta}; \tau),$$

where the summation is over all permutations over  $[n]$ . In view of this connection, the ANOPE constitutes another subsampled version of the Rank Lasso estimator.

Both ANOPE and the subsampling strategy adopted by Wang, Peng, et al. (2020) are incomplete  $U$ -statistics that use  $mn$  out of  $n(n - 1)/2$  terms to approximate the complete version  $L_n(\boldsymbol{\beta})$  in (3). All sampling mechanisms generating  $mn$  pairs of indices from  $\{(i, j) : 1 \leq i < j \leq n\}$  independently of the data  $\{\mathbf{x}_i, y_i\}_{i=1}^n$  produce unbiased estimates for  $\mathbb{E}L_n(\boldsymbol{\beta})$ . Among them, ANOPE has the minimum variance, according to Example 1 in Lee (2019, sec. 4.3.2). In addition, the first few  $m$  provide most variance reduction, as new pairs provide more independent information. When  $m$  is sufficiently large, additional pairs do not add much information. Let us illustrate these by a simulation study.

To compare the two different subsampling-based estimators, we generate  $\mathbf{x}_i$  from  $\mathcal{N}(\mathbf{0}, \mathbf{I}_p)$  and  $\varepsilon_i \sim \mathcal{N}(0, 1)$ . We take  $n = 100, p = 400$ , and  $\boldsymbol{\beta}_0 = (\sqrt{3}, \sqrt{3}, \sqrt{3}, 0, \dots, 0)^\top$ . We vary  $m$  from 1 to 20, and in addition we take into account the full sample regime where no subsampling is used. Note that same  $m$  yields the same number of pairwise losses in both estimators. For the purpose of comparison, we use the same error metrics as in Wang, Peng, et al. (2020), namely the  $\ell_1$  error,  $\ell_2$  error, the number of false positives and the number of false negatives. Their averages over 200 trials are plotted in Figure 1. Two crucial observations are worth mentioning. First, for a wide range of choices of  $m$ , the estimator based on subsampling permutations performs uniformly better. Second,  $m = 5$  seems to be a good choice for practical implementations to balance computational and statistical efficiency. Third, even with  $m = 1$ , approximately the same computation as penalized LAD, the efficiency of ANOPE is already very high, approximately  $(\frac{1.26}{1.39})^2 = 0.82$  in  $\ell_2$  error. Therefore, for the remaining experiments in this article, we take  $m = 5$ , though we are aware  $m$  depends on various factors in the simulation, including the tails of the design and noise.

### 3. Choice of the Regularization Parameter $\lambda$

In this section, we focus on the choice of the regularization parameter  $\lambda$  in (6). Let  $\mathbf{S}(\mathbf{y} - \mathbf{X}\boldsymbol{\beta})$  be the negative gradient of  $D(\mathbf{y} - \mathbf{X}\boldsymbol{\beta})^1$ , that is,

$$\mathbf{S}(\mathbf{y} - \mathbf{X}\boldsymbol{\beta}) = -\nabla D(\mathbf{y} - \mathbf{X}\boldsymbol{\beta}) = \frac{\sqrt{3}}{n+1} \mathbf{X}^\top [2\mathbf{r}(\boldsymbol{\beta}) - (n+1)], \tag{8}$$


with  $\mathbf{r}(\boldsymbol{\beta}) \in \mathbb{R}^n$  being the rank vector of the residuals  $\mathbf{y} - \mathbf{X}\boldsymbol{\beta}$ . The gradient at  $\boldsymbol{\beta} = \boldsymbol{\beta}_0$  plays an important role in the choice of the regularization parameter  $\lambda$ . Note that  $\mathbf{S}_n$  defined in the article has the following equivalent representation

$$\mathbf{S}_n = -\frac{2}{\sqrt{3}} \frac{n+1}{\sqrt{n(n-1)}} \frac{1}{\sqrt{n}} \mathbf{S}(\mathbf{y} - \mathbf{X}\boldsymbol{\beta}_0).$$

The recommended choice of  $\lambda$  is then given by

$$\lambda^* = c \cdot G_{\|\mathbf{S}_n\|_\infty}^{-1}(1 - \alpha_0)$$

<sup>1</sup>Since  $D(\mathbf{y} - \mathbf{X}\boldsymbol{\beta})$  is piecewise linear in  $\boldsymbol{\beta}$ , the negative gradient  $\mathbf{S}(\mathbf{y} - \mathbf{X}\boldsymbol{\beta})$  is defined everywhere excluding the linear boundaries.



**Figure 1.** Different error metrics for estimating  $\beta_0$  versus the sampling budget  $m$ . (a)  $\ell_1$  error, (b)  $\ell_2$  error, (c) false positives, and (d) false negatives. The results are based on the average of 200 simulations.

with  $c = 1.01$  and  $\alpha_0 = 0.1$ , where  $G_{\|\mathbf{S}_n\|_\infty}^{-1}(1 - \alpha_0)$  denotes the  $(1 - \alpha_0)$ -quantile of the distribution of  $\|\mathbf{S}_n\|_\infty$ . We now develop a basic understanding of the distribution of  $\mathbf{S}_n$ . It has been shown in Hettmansperger and McKean (2010, Theorem 3.5.2) that

$$\frac{1}{\sqrt{n}}\mathbf{S}(\mathbf{y} - \mathbf{X}\beta_0) \xrightarrow{d} \mathcal{N}(\mathbf{0}, \Sigma)$$

in the fixed  $p$ , large  $n$  regime. Here,  $\Sigma := \lim_n \mathbf{X}'\mathbf{X}/n$  is the limit of the covariance matrix of  $\mathbf{X}$ . This motivates another way to select the regularization parameter

$$\lambda^{\text{new}} = 1.01 \cdot \frac{2}{\sqrt{3}} \frac{n+1}{\sqrt{n}(n-1)} G_{\|\mathbf{v}\|_\infty}^{-1}(1 - \alpha_0),$$

where  $\mathbf{v} \sim \mathcal{N}(\mathbf{0}, \mathbf{X}'\mathbf{X}/n)$ , which can be simulated via multiplier bootstrap, that is,  $\mathbf{v} = n^{-1/2} \sum_{i=1}^n \eta_i \mathbf{X}_i$  with  $\eta_i \sim \mathcal{N}(0, 1)$ . The consistency of the multiplier bootstrap has been established (Chernozhukov, Chetverikov, and Kato 2014; Fan, Shao, and Zhou 2018). An advantage of this choice is it does not depend on the unknown  $\beta_0$  and can be computed before running Rank Lasso.

### 4. Bridging Rank Lasso and Other Pivotal Procedures

Thanks to the pivotal property of  $L_n$ 's subgradient function (see Wang, Peng, et al. 2020, sec. 2.2), tuning can be easily done via simulation without any knowledge of the error distribution. In particular, the selection of  $\lambda$  is not affected by the size of noise. In stark contrast, the optimal  $\lambda$  for Lasso

$$\min_{\beta \in \mathbb{R}^p} \left\{ \frac{1}{2n} \|\mathbf{X}\beta - \mathbf{y}\|_2^2 + \lambda \|\beta\|_1 \right\} \tag{9}$$

is of order  $n^{-1} \|\mathbf{X}^\top \boldsymbol{\varepsilon}\|_\infty$  that scales linearly in the standard deviation of  $\varepsilon_i$ 's.

We now take a closer look at the loss function  $L_n$  in (3) under a Gaussian model and then draw links between the Rank Lasso and two other popular approaches, square-root Lasso (Belloni, Chernozhukov, and Wang 2011) and scaled Lasso (Sun and Zhang 2012), with similar pivotal properties. Suppose that  $\{\mathbf{x}_i\}_{i=1}^n$  are iid copies of  $N(\mathbf{0}, \Sigma)$  and  $\boldsymbol{\varepsilon} \sim N(\mathbf{0}, \sigma^2 \mathbf{I}_n)$  is independent of  $\mathbf{X}$ . By (1), the population version of  $L_n$  is

$$L(\beta) = \mathbb{E}L_n(\beta) = \mathbb{E}|(y_1 - \mathbf{x}_1^\top \beta) - (y_2 - \mathbf{x}_2^\top \beta)|.$$

Since

$$y - X\beta \sim N(\mathbf{0}, [\sigma^2 + (\beta - \beta_0)^\top \Sigma (\beta - \beta_0)] \mathbf{I}_n) \quad (10)$$

has iid centered Gaussian entries, we have

$$\begin{aligned} L(\beta) &= \sqrt{\frac{2}{\pi}} \mathbb{E}^{1/2} |(y_1 - \mathbf{x}_1^\top \beta) - (y_2 - \mathbf{x}_2^\top \beta)|^2 \\ &= \frac{2}{\sqrt{\pi}} \mathbb{E}^{1/2} |y_1 - \mathbf{x}_1^\top \beta|^2 = \frac{2}{\sqrt{\pi n}} \mathbb{E}^{1/2} \|y - X\beta\|_2^2. \end{aligned} \quad (11)$$

Then

$$L(\beta) + \lambda \|\beta\|_1 = \frac{2}{\sqrt{\pi}} \left( n^{-1} \mathbb{E} \|y - X\beta\|_2^2 \right)^{1/2} + \lambda \|\beta\|_1.$$

The right-hand side can be viewed as a population version of the penalized loss in square-root Lasso (Belloni, Chernozhukov, and Wang 2011).

On the other hand, (10) and (11) lead to

$$\begin{aligned} L(\beta) &= \frac{2}{\sqrt{\pi}} \sqrt{\sigma^2 + (\beta - \beta_0)^\top \Sigma (\beta - \beta_0)} \\ &= \frac{2\sigma}{\sqrt{\pi}} [1 + \sigma^{-2} \mathbb{E} |\mathbf{x}_1^\top (\beta - \beta_0)|^2]^{1/2}. \end{aligned}$$

The term

$$\mathbb{E} |\mathbf{x}_1^\top (\beta - \beta_0)|^2 = \mathbb{E} |y_1 - \mathbf{x}_1^\top \beta|^2 - \mathbb{E} |y_1 - \mathbf{x}_1^\top \beta_0|^2$$

is the excess risk of  $\beta$ . When  $\mathbb{E} |\mathbf{x}_1^\top (\beta - \beta_0)|^2 \ll \sigma^2$ , we use Taylor expansion to derive

$$\begin{aligned} L(\beta) &\approx \frac{2\sigma}{\sqrt{\pi}} \left( 1 + \frac{1}{2\sigma^2} \mathbb{E} |\mathbf{x}_1^\top (\beta - \beta_0)|^2 \right) \\ &= \frac{2\sigma}{\sqrt{\pi}} \left( 1 + \frac{\mathbb{E} |y_1 - \mathbf{x}_1^\top \beta|^2 - \mathbb{E} |y_1 - \mathbf{x}_1^\top \beta_0|^2}{2\sigma^2} \right) \\ &= \frac{\sigma}{\sqrt{\pi}} \left( 1 + \frac{\mathbb{E} |y_1 - \mathbf{x}_1^\top \beta|^2}{\sigma^2} \right). \end{aligned}$$

Therefore,

$$L(\beta) + \lambda \|\beta\|_1 \approx \frac{1}{\sqrt{\pi}} \left( \sigma + \frac{1}{n\sigma} \mathbb{E} \|y - X\beta\|_2^2 \right) + \lambda \|\beta\|_1.$$

This time the right-hand side becomes a population version of the penalized loss in scaled Lasso (Sun and Zhang 2012).

### 5. Inference Based on Rank Lasso

Here, we explore the possibility of conducting inference (e.g., constructing confidence intervals for coordinates in  $\beta_0$ ) using the Rank Lasso estimator. As in the standard Lasso estimator, the Rank Lasso estimator  $\hat{\beta}$  is no longer unbiased for  $\beta_0$  due to the  $\ell_1$  regularizer. Inspired by the de-biased Lasso estimator (Javanmard and Montanari 2014; Van de Geer et al. 2014; Zhang and Zhang 2014), a natural approach to inference is to remove the bias in  $\hat{\beta}$  and construct an asymptotically unbiased estimator  $\hat{\beta}^d$ , whose distribution is easy to characterize.

In what follows, we sketch a heuristic argument that would ultimately lead us toward a de-biased estimator. In view of the

definition (6),  $\hat{\beta}$  satisfies the first-order optimality condition of (6)

$$-\frac{2(n+1)}{\sqrt{3n(n-1)}} \mathbf{S}(y - X\hat{\beta}) + \lambda \partial \|\hat{\beta}\|_1 = \mathbf{0}. \quad (12)$$

Here,  $\partial \|\hat{\beta}\|_1$  denotes a subgradient of  $\|\beta\|_1$  at  $\beta = \hat{\beta}$ ,  $\mathbf{S}(y - X\beta)$  is the negative gradient of  $D(y - X\beta)$  defined in (8). In view of the asymptotic theory for rank based estimators in the low dimensional setting,  $\mathbf{S}(y - X\beta)$  admits a linear approximation (Hettmansperger and McKean 2010, Theorem A.3.1), that is,

$$\begin{aligned} \frac{1}{\sqrt{n}} \mathbf{S}(y - X\beta) &\approx \frac{1}{\sqrt{n}} \mathbf{S}(y - X\beta_0) \\ &\quad - \sqrt{12} \int f^2(u) du \Sigma \cdot \sqrt{n} (\beta - \beta_0), \end{aligned} \quad (13)$$

where  $\Sigma$  is the sample covariance of  $X$  and  $f(u)$  denotes the density function of the error  $\epsilon$ . Combine (12) and (13) to reach<sup>2</sup>

$$\begin{aligned} \sqrt{n} \left( \hat{\beta} + \Sigma^{-1} \frac{1}{\int f^2(u) du} \frac{n-1}{4(n+1)} \lambda \partial \|\hat{\beta}\|_1 - \beta_0 \right) \\ \approx \frac{1}{\sqrt{12} \int f^2(u) du} \Sigma^{-1} \frac{1}{\sqrt{n}} \mathbf{S}(y - X\beta_0). \end{aligned}$$

Use the optimality condition (12) and the definition of  $\mathbf{S}(y - X\beta)$  to arrive at

$$\begin{aligned} \sqrt{n} \left( \hat{\beta} + \frac{1}{\int f^2(u) du} \frac{1}{2n(n+1)} \Sigma^{-1} \mathbf{X}^\top (2r(\hat{\beta}) - (n+1)) - \beta_0 \right) \\ \approx \frac{1}{\sqrt{12} \int f^2(u) du} \Sigma^{-1} \frac{1}{\sqrt{n}} \mathbf{S}(y - X\beta_0) \\ = \frac{1}{\int f^2(u) du} \frac{1}{2\sqrt{n}(n+1)} \Sigma^{-1} \mathbf{X}^\top (2r(\beta_0) - (n+1)) =: \Xi. \end{aligned}$$

Clearly, the right-hand side  $\Xi$  has zero mean and is asymptotically normal. In fact,  $\Xi$  is closely related to  $\mathbf{S}_n$  defined in the article:

$$\frac{1}{\sqrt{n}} \Xi = \frac{1}{4 \int f^2(u) du} \Sigma^{-1} \mathbf{S}_n. \quad (14)$$

This motivates the construction of the following de-biased estimator<sup>3</sup>

$$\hat{\beta}^d := \hat{\beta} + \frac{1}{\int f^2(u) du} \frac{1}{2n(n+1)} \Sigma^{-1} \mathbf{X}^\top (2r(\hat{\beta}) - (n+1)). \quad (15)$$

In addition, for each  $1 \leq j \leq p$ , a valid  $(1 - \alpha)$ -confidence interval of  $\beta_{0j}$  is given by

$$[\hat{\beta}_j^d - G_{\Xi_j/\sqrt{n}}^{-1}(1 - \alpha/2), \hat{\beta}_j^d - G_{\Xi_j/\sqrt{n}}^{-1}(1 - \alpha/2)],$$

where  $G_{\Xi_j/\sqrt{n}}^{-1}(1 - \alpha/2)$  denotes the  $(1 - \alpha/2)$ -quantile of the distribution of  $\Xi_j/\sqrt{n}$ .

Below, we empirically demonstrate the validity of the de-biased estimator  $\hat{\beta}^d$ . The experimental setup is similar to that

<sup>2</sup>For simplicity, here we assume the sample covariance is invertible (which requires  $n > p$ ), otherwise an estimator of the inverse covariance is needed as in Javanmard and Montanari (2014).

<sup>3</sup>A consistent estimator of  $\int f^2(u) du$  is needed to make this fully practical.



**Table 1.** Average coverage probabilities for 90% confidence intervals.

|        | Coverage ( $\mathcal{S}_0$ ) | Coverage ( $\mathcal{S}_0^c$ ) | Coverage ( $[p]$ ) |
|--------|------------------------------|--------------------------------|--------------------|
| Normal | 89.67%                       | 90.71%                         | 90.70%             |
| Cauchy | 88.50%                       | 90.43%                         | 90.42%             |

in Section 2 except that we set  $n = 600$  and  $p = 400$ . For any set  $\mathcal{S} \subseteq [p]$ , we define the coverage probability of  $\hat{\beta}^d$  on  $\mathcal{S}$  to be

$$\text{Coverage}(\mathcal{S}) = \frac{1}{|\mathcal{S}|} \sum_{j \in \mathcal{S}} \mathbb{1} \left\{ \beta_{0j} \in [\hat{\beta}_j^d - G_{\Xi_j/\sqrt{n}}^{-1}(1 - \alpha/2), \hat{\beta}_j^d + G_{\Xi_j/\sqrt{n}}^{-1}(1 - \alpha/2)] \right\}.$$

In particular, we focus on the coverage probability on important variables  $\mathcal{S}_0 := \{j \in [p] \mid \beta_{0j} \neq 0\}$ , unimportant variables  $\mathcal{S}_0^c$  and all the variables  $\Omega = [p]$ . Table 1 reports the averaged results over 200 Monte Carlo simulations for  $1 - \alpha = 0.9$ . As can be seen from Table 1, the average coverage probabilities are quite close to the nominal level 90%. While the de-biasing approach gives correct coverage, the confidence intervals are much wider than the oracle estimator, for both active and inactive components. For example, for inactive components, any intervals, however small, give 100% coverage, as long as they contain the origin. How to construct confidence intervals taking more into account of the lengths of the intervals, in addition to the correct coverage? This requires more effective use of sparsity.

### 6. Regularization Under Strongly Dependent Covariates

The main theorem is established under restricted eigenvalue condition in Wang, Peng, et al. (2020). This requires the weakly dependent covariates, which usually do not hold in high-dimensional setting, as high-dimensional covariates often measures similar things (e.g., economic health, financial returns, gene expressions). The strongly dependent covariates are often modeled through common factors and the factor adjustments are needed in the regularization (Fan, Ke, and Wang 2020).

Suppose that  $\{x_i\}_{i=1}^n$  are generated from the approximate factor model

$$x_i = Bf_i + u_i, \quad i \in [n], \tag{16}$$

where  $B \in \mathbb{R}^{p \times K}$  is a loading matrix,  $f_i \in \mathbb{R}^K$  gives the latent factors, and  $u_i \in \mathbb{R}^p$  records idiosyncratic components which are weak dependent. Factor-adjusted regularized model selection (FarmSelect) (Fan, Ke, and Wang 2020) decorrelates the covariates as follows. Note that

$$y_i = x_i^\top \beta_0 + \varepsilon_i = f_i^\top B^\top \beta_0 - u_i^\top \beta_0 + \varepsilon_i. \tag{17}$$

If  $\{f_i, u_i\}_{i=1}^n$  are observable, then we can perform Rank Lasso using  $(f_i, u_i)$  rather than  $x_i$ :


$$(\hat{\gamma}, \hat{\beta}) \in \underset{\gamma \in \mathbb{R}^K, \beta \in \mathbb{R}^p}{\operatorname{argmin}} \left\{ \frac{1}{n(n-1)} \sum_{i \neq j} |(y_i - f_i^\top \gamma - u_i^\top \beta) - (y_j - f_j^\top \gamma - u_j^\top \beta)| + \lambda \|\beta\|_1 \right\}, \tag{18}$$

and then use  $\hat{\beta}$  as the estimate for  $\beta_0$ . Here, we only enforce the sparsity of  $\beta$ . Regression using the factors and idiosyncratic components facilitates model selection as the new covariates have weaker correlations. When the factors are not observable, we replace  $\{f_i, u_i\}_{i=1}^n$  in (18) by their empirical estimates from  $\{x_i\}_{i=1}^n$  using the principal component analysis.

We now demonstrate the efficacy of the procedure above on synthetic data. Consider the linear model (1) with  $n = 100$ ,  $p = 400$ ,  $\beta_0 = (2, 2, 2, 0, \dots, 0)^\top \in \mathbb{R}^p$  and  $\varepsilon_i \sim N(0, 1)$ . Let  $x_i \sim N(\mathbf{0}, \rho \mathbf{1}\mathbf{1}^\top + (1 - \rho)\mathbf{I})$  with  $\rho = 0.2$ . This is equivalent to the factor model (16) with  $K = 1$ ,  $B = \sqrt{\rho}\mathbf{1} \in \mathbb{R}^p$ ,  $f_i \sim N(0, 1)$ , and  $u_i \sim N(\mathbf{0}, (1 - \rho)\mathbf{I})$  independent of  $f_i$ . We compare the Rank Lasso estimator (2) and its decorrelated version (18) with  $f_i$  and  $u_i$  estimated using principal component analysis, see Section 3.1 in Fan, Ke, and Wang (2020). Throughout the experiment, we choose the penalty parameters using the simulation method (Wang, Peng, et al. 2020) with 500 replicates, fix  $\alpha_0 = 0.1$  and vary  $c$  from 1 to 2.2 by 0.01. To speedup computation, we subsample  $5n$  pairs in  $\{(i, j)\}_{1 \leq i < j \leq n}$  with replacement to approximate the  $U$ -statistics. Figure 2 shows the model selection errors, which are averaged over 200 independent runs.

The selection error is the total number of false positives and false negatives, that is, the symmetric difference between the selected set of variables and the true one. Since the average selection error of Rank Lasso is bounded away from zero, it cannot consistently identify the true set. Fortunately, decorrelation helps select the true model. When both methods achieve their minimum selection error ( $c \approx 1.6$  for Rank Lasso and  $c \approx 1.15$  for the decorrelated one), the  $\ell_2$  error of the Rank Lasso (1.84) is higher than that of its decorrelated version (1.47). The figure also shows that the regularization parameter  $c$  or equivalently  $\lambda$  is sensitive to the regularization.

If the covariates are correlated,  $\ell_1$ -regularized sparse regression needs a large penalty parameter to return a set of variables close to the true one. That inevitably induces high bias and may jeopardize the estimation accuracy. While model selection and parameter estimation can be at odds with each other, the decorrelation step before regression provides a reconciliation.



**Figure 2.** Rank Lasso and decorrelation: selection errors (false positives + false negatives).

## 7. Further Comments

There is a large literature on robust high-dimension regression based on Huber type of loss with diverging tuning parameter  $\tau$  (see, e.g., Catoni 2012; Devroye et al. 2016; Fan, Li, and Wang 2017; Fan, Wang, and Zhu 2020; Sun, Zhou, and Fan 2020). Wang, Zheng, et al. (2020) propose a data-driven method of the robustification parameter. The pros and cons of using Rank Lasso or adaptive Huber regression cycle back to those in the low-dimensional setting. Rank Lasso requires no moment conditions, no tuning parameters in the loss, but less efficient where error is normal and requires the error density bounded away from zero at original. This also explains that the minimax result of non- $\sqrt{n}$ -consistency of Sun, Zhou, and Fan (2020) bears no contradiction with the result in Wang, Peng, et al. (2020).

The current article (Wang, Peng, et al. 2020) analyze theoretical properties for the Rank Lasso under a homoscedastic model (1) where  $\{\varepsilon_i\}_{i=1}^n$  are iid given  $\{\mathbf{x}_i\}_{i=1}^n$ . For such model, both the conditional median and the conditional mean of  $y_i$  given  $\mathbf{x}_i$  are affine functions of  $\mathbf{x}_i$ , and they differ only by the intercept. Based on the observation, the  $\ell_1$ -regularized least absolute deviations

$$\min_{\gamma \in \mathbb{R}, \boldsymbol{\beta} \in \mathbb{R}^p} \left\{ \frac{1}{n} \sum_{i=1}^n |y_i - (\gamma + \mathbf{x}_i^\top \boldsymbol{\beta})| + \lambda \|\boldsymbol{\beta}\|_1 \right\} \quad (19)$$

should also serve the needs. This is computationally more attractive as we get rid of the  $U$ -statistic. The same intuition also holds for general quantile regression. Results on robustness and tuning are available (see Belloni and Chernozhukov 2011). To better demonstrate the efficacy of Rank Lasso, one could consider, for example, a class of heteroscedastic model where  $\{(\mathbf{x}_i, \varepsilon_i)\}_{i=1}^n$  are iid but the conditional distribution of  $\varepsilon_i$  given  $\mathbf{x}_i = \mathbf{x}$  varies with  $\mathbf{x}$ . Quantile regression breaks down in this scenario. Thanks to the symmetrization step for constructing the loss (3), the Rank Lasso is still expected to enjoy nice theoretical guarantees.

In addition, it would be nice to see how to go beyond linear regression. While Rank Lasso is a convex optimization problem, things are complicated for generalized linear models such as the logistic model. It seems challenging to design a loss function, rank-based or not, that still possesses robustness and pivotal properties for nonlinear problems.

## Funding

The research was supported by NSF grants DMS-1662139, DMS-1712591, NIH grant 2R01-GM072611-16, and ONR grant N00014-19-1-2120.

## References

- Bassett, G., and Koenker, R. (1978), "Asymptotic Theory of Least Absolute Error Regression," *Journal of the American Statistical Association*, 73, 618–622. [1721]
- Belloni, A., and Chernozhukov, V. (2011), " $\ell_1$ -Penalized Quantile Regression in High-Dimensional Sparse Models," *The Annals of Statistics*, 39, 82–130. [1721,1725]
- Belloni, A., Chernozhukov, V., and Wang, L. (2011), "Square-Root Lasso: Pivotal Recovery of Sparse Signals via Conic Programming," *Biometrika*, 98, 791–806. [1722,1723]
- Catoni, O. (2012), "Challenging the Empirical Mean and Empirical Variance: A Deviation Study," *Annales de l'IHP Probabilités et Statistiques*, 48, 1148–1185. [1725]
- Chernozhukov, V., Chetverikov, D., and Kato, K. (2014), "Gaussian Approximation of Suprema of Empirical Processes," *The Annals of Statistics*, 42, 1564–1597. [1722]
- Devroye, L., Lerasle, M., Lugosi, G., and Oliveira, R. I. (2016), "Sub-Gaussian Mean Estimators," *The Annals of Statistics*, 44, 2695–2725. [1725]
- Fan, J., Fan, Y., and Barut, E. (2014), "Adaptive Robust Variable Selection," *The Annals of Statistics*, 42, 324. [1721]
- Fan, J., Ke, Y., and Wang, K. (2020), "Factor-Adjusted Regularized Model Selection," *Journal of Econometrics*, 2020. [1724]
- Fan, J., Li, Q., and Wang, Y. (2017), "Estimation of High Dimensional Mean Regression in the Absence of Symmetry and Light Tail Assumptions," *Journal of the Royal Statistical Society, Series B*, 79, 247. [1725]
- Fan, J., Shao, Q.-M., and Zhou, W.-X. (2018), "Are Discoveries Spurious? Distributions of Maximum Spurious Correlations and Their Applications," *The Annals of Statistics*, 46, 989. [1722]
- Fan, J., Wang, W., and Zhu, Z. (2020), "A Shrinkage Principle for Heavy-Tailed Data: High-Dimensional Robust Low-Rank Matrix Recovery," *The Annals of Statistics* (to appear). [1725]
- Hettmansperger, T. P., and McKean, J. W. (1978), "Statistical Inference Based on Ranks," *Psychometrika*, 43, 69–79. [1720]
- (2010), *Robust Nonparametric Statistical Methods*, Boca Raton, FL: CRC Press. [1720,1722,1723]
- Hodges, J. L., Jr., and Lehmann, E. L. (1963), "Estimates of Location Based on Rank Tests," *The Annals of Mathematical Statistics*, 34, 598–611. [1720]
- Javanmard, A., and Montanari, A. (2014), "Confidence Intervals and Hypothesis Testing for High-Dimensional Regression," *The Journal of Machine Learning Research*, 15, 2869–2909. [1723]
- Lee, A. J. (2019), *U-Statistics: Theory and Practice*, New York: Routledge. [1721]
- Sun, Q., Zhou, W.-X., and Fan, J. (2020), "Adaptive Huber Regression," *Journal of the American Statistical Association*, 115, 254–265. [1725]
- Sun, T., and Zhang, C.-H. (2012), "Scaled Sparse Linear Regression," *Biometrika*, 99, 879–898. [1722,1723]
- Van de Geer, S., Bühlmann, P., Ritov, Y., and Dezeure, R. (2014), "On Asymptotically Optimal Confidence Regions and Tests for High-Dimensional Models," *The Annals of Statistics*, 42, 1166–1202. [1723]
- Wang, L., Peng, B., Bradic, J., Li, R., and Wu, Y. (2020), "A Tuning-Free Robust and Efficient Approach to High-Dimensional Regression," *Journal of the American Statistical Association*. [1720,1721,1722,1724,1725]
- Wang, L., Zheng, C., Zhou, W., and Zhou, W.-X. (2020), "A New Principle for Tuning-Free Huber Regression," *Statistica Sinica*. [1725]
- Zhang, C.-H., and Zhang, S. S. (2014), "Confidence Intervals for Low Dimensional Parameters in High Dimensional Linear Models," *Journal of the Royal Statistical Society, Series B*, 76, 217–242. [1723]