

# Efficient Estimation and Inferences for Varying-Coefficient Models

Zongwu CAI, Jianqing FAN, and Runze LI

This article deals with statistical inferences based on the varying-coefficient models proposed by Hastie and Tibshirani. Local polynomial regression techniques are used to estimate coefficient functions, and the asymptotic normality of the resulting estimators is established. The standard error formulas for estimated coefficients are derived and are empirically tested. A goodness-of-fit test technique, based on a nonparametric maximum likelihood ratio type of test, is also proposed to detect whether certain coefficient functions in a varying-coefficient model are constant or whether any covariates are statistically significant in the model. The null distribution of the test is estimated by a conditional bootstrap method. Our estimation techniques involve solving hundreds of local likelihood equations. To reduce the computational burden, a one-step Newton–Raphson estimator is proposed and implemented. The resulting one-step procedure is shown to save computational cost on an order of tens with no deterioration in performance, both asymptotically and empirically. Both simulated and real data examples are used to illustrate our proposed methodology.

**KEY WORDS:** Asymptotic normality; Bootstrap; Generalized linear models; Goodness-of-fit; Local polynomial fitting; One-step procedure.

## 1. INTRODUCTION

Generalized linear models are based on two fundamental assumptions: the conditional distributions belong to an exponential family and a known transform of the underlying regression function is linear. In recent years, various attempts have been made to relax these model assumptions and hence widen their applicability, because a wrong model on the regression function can lead to excessive modeling biases and erroneous conclusions. Of importance is the varying-coefficient models, proposed by Hastie and Tibshirani (1993), which widen the scope of applications by allowing regression coefficients to depend on certain covariates.

A varying-coefficient model has the form

$$\eta(\mathbf{u}, \mathbf{x}) = g\{m(\mathbf{u}, \mathbf{x})\} = \sum_{j=1}^p a_j(\mathbf{u})x_j \quad (1)$$

for some given link function  $g(\cdot)$ , where  $\mathbf{x} = (x_1, \dots, x_p)^T$  and  $m(\mathbf{u}, \mathbf{x})$  is the mean regression function of the response variable  $Y$  given the covariates  $\mathbf{U} = \mathbf{u}$  and  $\mathbf{X} = \mathbf{x}$  with  $\mathbf{X} = (X_1, \dots, X_p)^T$ . Clearly, (1) includes both the parametric generalized linear model (McCullagh and Nelder 1989) and the generalized partially linear model (Carroll, Fan, Gijbels, and Wand 1997; Chen 1988; Green and Silverman 1994; Speckman 1988).

A motivation of this study comes from an analysis of environmental data, consisting of weekly measurements of pollutants and other environmental factors, collected in Hong Kong from January 1, 1994 to December 31, 1995 (courtesy of T. S. Lau). Of interest is to examine the association between the levels of pollutants and the total number of weekly hospital admissions for circulatory and res-

piratory problems. It is natural to allow the association to change over time (see Fig. 3 later); such a problem can be tackled by using (1) as follows. The log-link is used,  $\mathbf{U}$  is the time covariate, and  $\mathbf{X}$  denotes the levels of pollutants. The conditional distribution of the number of weekly hospital admissions given the covariates is modeled as a Poisson distribution with the mean function given by (1). In another context, one is interested in studying how the variables such as burn area and gender affect survival probabilities for different age of burn victims. Detailed analysis of these two datasets is reported in Section 3.

In the least-squares setting, (1) with the identity link was introduced by Cleveland, Grosse, and Shyu (1992) and extended by Hastie and Tibshirani (1993) to various aspects. Recently, some new development has been made to (1). Kauermann and Tutz (1999) proposed a graphical procedure to diagnose the discrepancy between the parametric model and the smoothing alternative by using the local likelihood smoothing. Furthermore, a two-step estimation procedure was proposed by Fan and Zhang (1999) to deal with the situations where coefficient functions admit different degrees of smoothness. An advantage of (1) is that by allowing the coefficients  $\{a_j(\cdot)\}$  to depend on  $\mathbf{U}$ , the modeling bias can be reduced significantly, and the “curse of dimensionality” is avoided.

Varying-coefficient models are a simple and useful extension of classical generalized linear models. This extension admits simple interpretation. The models are particularly appealing in longitudinal studies, where they allow one to explore the extent to which covariates affect responses changing over time. Brumback and Rice (1998), Fan and Zhang (2000), and Hoover, Rice, Wu, and Yang (1998) have given details on novel applications of the varying-coefficient models to longitudinal data. Chen and Tsay (1993) have explored nonlinear time series applications, and Cai, Fan, and Yao (1998) have provided statistical inferences based on the functional-coefficient autore-

Zongwu Cai is Assistant Professor, Department of Mathematics, University of North Carolina, Charlotte, NC 28223 (E-mail: [zcaai@unc.edu](mailto:zcaai@unc.edu)). Jianqing Fan is Professor, Department of Statistics, University of California, Los Angeles, CA 90095 (E-mail: [jfan@math.ucla.edu](mailto:jfan@math.ucla.edu)). Runze Li is Graduate Student, Department of Statistics, University of North Carolina, Chapel Hill, NC 27599 (E-mail: [lirz@email.unc.edu](mailto:lirz@email.unc.edu)). Fan’s research was partially supported by National Science Foundation grant DMS-9803200. The authors thank the editor, the associate editor, and two referees for their constructive and detailed suggestions that led to significant improvements in the presentation of the article.

gressive models. Cai et al. (1998) also gave an extensive study on the advantages of the varying-coefficient model over the parametric model based on the predictive utility. For applications in finance, econometrics and environmental study, we refer to the unpublished papers by Hong and Lee (1999) and Lee and Ullah (1999), and the article by Cai and Tiwari (1999).

Estimation of the coefficient functions in (1) is obtained by using local smoothing techniques. By localizing data around  $\mathbf{u}$ , (1) is approximately a generalized linear model. One can find its local maximum likelihood estimate (MLE) using an iterative algorithm. Note that the local MLE for the varying-coefficient model is indeed solving the local likelihood equations. Thus our local likelihood method can be regarded as a special case of the general local estimation equation method proposed by Carroll, Ruppert, and Welsh (1998). Hence the bandwidth involved can be selected by the empirical bias method proposed in that article. To obtain the estimated coefficient functions, we need to solve hundreds of local maximum likelihood problems. The computation can be expensive, depending on the convergence criterion. Computational burden becomes even more severe when a cross-validation method is used to select a smoothing parameter. To reduce computational costs, we propose a one-step local MLE. The idea is not novel, because it was first used by Bickel (1975) in the parametric setting, but implementations and insights are. We show that computational costs can be reduced significantly and demonstrate that the resulting one-step estimator is as efficient, both asymptotically and empirically, as the fully iterative MLE.

Associated with inferences on the varying-coefficient models are the standard errors of the estimated coefficient functions. We derive consistent estimates. Our simulation studies show that the estimated standard errors are very accurate for most applications. Another important issue arises regarding whether some of the coefficient functions in (1) are actually varying, or whether some of covariates are statistically significant. We propose a nonparametric maximum likelihood ratio test and estimate its null distribution using a conditional bootstrap method. Our simulation shows that the resulting testing procedure performs well.

One of our goals is to efficiently estimate the coefficient functions  $\{a_j(\cdot)\}$  in (1) by using a nonparametric method. Our methods are directly applicable to situations in which one cannot fully specify the conditional log-likelihood function  $l(v, y)$ , but can model the relationship between the mean and variance by  $\text{var}(Y|U = \mathbf{u}, \mathbf{X} = \mathbf{x}) = \sigma^2 V\{m(\mathbf{u}, \mathbf{x})\}$  for a known variance function  $V(\cdot)$  and unknown  $\sigma$ . In this case, one need only replace the log-likelihood function  $l(v, y)$  by the quasi-likelihood function  $Q(\cdot, \cdot)$ , defined by  $(\partial/\partial\mu)Q(\mu, y) = (y - \mu)/[V(\mu)]$ . We assume throughout that the conditional log-likelihood function  $l(v, y)$  is known and linear in  $y$  for fixed  $v$ . This assumption is satisfied for the canonical exponential family, which is the focus of this article.

The article is organized as follows. In Section 2 we discuss estimation methods and inference tools, and presents some asymptotic properties of the one-step and local MLEs.

In particular, formulas for consistent standard errors of the estimated coefficient functions are derived, a nonparametric maximum likelihood ratio test is proposed, and strategies are given for the implementation of a one-step estimator. In Section 3 we study some finite-sample properties of the one-step and local MLEs using two simulated examples. We illustrate our methodology through analysis of the aforementioned environmental and survival datasets. Finally, we give technical proofs in the Appendix.

## 2. MODELING PROCEDURES

For simplicity, we consider only the case in which  $\mathbf{u}$  is one-dimensional. Extension to multivariate  $\mathbf{u}$  involves no fundamentally new ideas. However, implementations with  $\mathbf{u}$  having more than two dimensions may have some difficulties due to the ‘‘curse of dimensionality.’’

### 2.1 Local Maximum Likelihood Estimate

We use a local linear modeling scheme, although general local polynomial methods are also applicable. The local linear fittings have several nice properties, such as high statistical efficiency (in an asymptotic minimax sense), design adaptation (Fan 1993), and good boundary behavior (Fan and Gijbels 1996; Ruppert and Wand 1994). Suppose that  $a_j(\cdot)$  has a continuous second derivative. For each given point  $u_0$ , we approximate  $a_j(u)$  locally by a linear function  $a_j(u) \approx a_j + b_j(u - u_0)$  for  $u$  in a neighborhood of  $u_0$ . Based on a random sample  $\{(U_i, \mathbf{X}_i, Y_i)\}_{i=1}^n$ , we use the following local likelihood method to estimate the coefficient functions:

$$l_n(\mathbf{a}, \mathbf{b}) = \frac{1}{n} \sum_{i=1}^n l \left[ g^{-1} \left\{ \sum_{j=1}^p (a_j + b_j(U_i - u_0)) X_{ij} \right\}, Y_i \right] \times K_h(U_i - u_0), \quad (2)$$

where  $K_h(\cdot) = K(\cdot/h)/h$ ,  $K(\cdot)$  is a kernel function,  $h = h_n > 0$  is a bandwidth,  $\mathbf{a} = (a_1, \dots, a_p)^T$ , and  $\mathbf{b} = (b_1, \dots, b_p)^T$ . Note that  $a_j$  and  $b_j$  are dependent on  $u_0$ , as is  $l_n(\cdot, \cdot)$ . Maximizing the local likelihood function  $l_n(\mathbf{a}, \mathbf{b})$  gives estimates  $\hat{\mathbf{a}}(u_0)$  and  $\hat{\mathbf{b}}(u_0)$ . The components in  $\hat{\mathbf{a}}(u_0)$  give an estimate of  $a_1(u_0), \dots, a_p(u_0)$ . For simplicity of notation, we denote  $\boldsymbol{\beta} = \boldsymbol{\beta}(u_0) = (a_1, \dots, a_p, b_1, \dots, b_p)^T$  and write the local likelihood function (2) as  $l_n(\boldsymbol{\beta})$ . Likewise, the local MLE is denoted by  $\hat{\boldsymbol{\beta}}_{\text{MLE}} = \hat{\boldsymbol{\beta}}_{\text{MLE}}(u_0)$ .

### 2.2 One-Step Local Maximum Likelihood Estimate

Computing the local MLE can be costly. This is especially true for varying-coefficient models. To obtain the estimated functions  $\{\hat{a}_j(\cdot)\}$ , one needs to maximize the local likelihood (2) for usually hundreds of distinct values of  $u_0$ , with each maximization requiring an iterative algorithm. Moreover, the computational expense further increases with the number of covariates  $p$ . To ameliorate this expense, we propose replacing the iterative local MLE by the one-step Newton–Raphson estimator, which has been frequently used in parametric models (Bickel 1975; Lehmann 1983). We prove Theorem 2 that the one-step local MLE does not

lose any statistical efficiency provided that the initial estimator is good enough.

Let  $l'_n(\beta)$  and  $l''_n(\beta)$  be the gradient and the Hessian matrix of the local log-likelihood  $l_n(\beta)$ . Given an initial estimator  $\hat{\beta}_0 = \hat{\beta}_0(u_0) = (\hat{a}(u_0)^T, \hat{b}(u_0)^T)^T$ , one step of the Newton-Raphson algorithm produces the updated estimator,

$$\hat{\beta}_{OS} = \hat{\beta}_0 - \{l''_n(\hat{\beta}_0)\}^{-1} l'_n(\hat{\beta}_0), \quad (3)$$

thus featuring the computational expediency of least-squares local polynomial fitting. In univariate generalized linear models, Fan and Chen (1999) carefully studied properties of the local one-step estimator. In that setting, the least-squares estimate serves a natural candidate as an initial estimator; however, in the multivariate setting, it is not clear how an initial estimator can be constructed.




Figure 1. Simulation Results for Example 1 With Sample Size 400. (a) The boxplots for the ratios of RASE of the one-step and two-step local likelihood approaches to that of the local MLE of  $\mathbf{a}(u)$ , using bandwidths (from left to right)  $h = .10, .20$ , and  $.40$ . (b), (c), and (d) Typical estimates of  $a_0(u)$ ,  $a_1(u)$ , and  $a_2(u)$ , with bandwidth  $h = .2$ . The solid curve represents the true function; the dashed curves (from shortest to longest dash) are the one-step, two-step and local MLE. (e) The estimated densities of  $T$  for unconditional null distributions (thick curves) and for conditional null distributions (thin curves). (f) The power functions of the test statistic  $T$ .

Note that  $l''_n(\hat{\beta}_0)$  can be nearly singular for certain  $u_0$ , due to possible data sparsity in certain local regions, or when the bandwidth is too small. Fan and Chen (1999) and Seifert and Gasser (1996) have explored the use of the ridge regression as an approach to handling such problems in the univariate setting. We extend their ideas in Section 3.

### 2.3 Sampling Properties

We now derive the asymptotic distributions of the local MLE  $\hat{\beta}_{MLE}$  and the one-step estimator  $\hat{\beta}_{OS}$ . We demonstrate that the one-step estimator performs as well as the local MLE, as long as the initial estimator  $\hat{\beta}_0$  is reasonably accurate.

Define  $\mu_k = \int u^k K(u) du$  and  $\nu_k = \int u^k K^2(u) du$ . Let  $\mathbf{H} = \text{diag}(1, h) \otimes \mathbf{I}_p$ , with  $\otimes$  denoting the Kronecker product. Let  $f_U(\cdot)$  denote the marginal density of  $U$ ,

$$\Gamma(u) = E\{\rho(U, \mathbf{X})\mathbf{X}\mathbf{X}^T | U = u\} \quad (4)$$

and

$$\rho(u, \mathbf{x}) = [g_1\{m(u, \mathbf{x})\}]^2 \text{var}\{Y | U = u, \mathbf{X} = \mathbf{x}\}, \quad (5)$$

with  $g_1(s) = g'_0(s)/g'(s)$  and  $g_0(\cdot)$  as the canonical link. Note that  $\rho(u, \mathbf{x}) = V\{m(u, \mathbf{x})\}$  for the canonical exponential family with the canonical link function. The asymptotic properties of  $\hat{\beta}_{MLE}$  and  $\hat{\beta}_{OS}$  are described in the following theorems, and conditions and proofs are given in the Appendix.

*Theorem 1.* Suppose that conditions C1–C7 in the Appendix hold and that  $h = h_n \rightarrow 0$  and  $nh \rightarrow \infty$  as  $n \rightarrow \infty$ . Then

$$\begin{aligned} \sqrt{nh} \left[ \mathbf{H}\{\hat{\beta}_{MLE}(u_0) - \beta(u_0)\} - \frac{h^2}{2(\mu_2 - \mu_1^2)} \right. \\ \times \begin{pmatrix} (\mu_2^2 - \mu_1\mu_3)\mathbf{a}''(u_0) \\ (\mu_3 - \mu_1\mu_2)\mathbf{a}''(u_0) \end{pmatrix} \\ \left. + o_p(h^2) \right] \xrightarrow{\mathcal{D}} \mathbf{N}(\mathbf{0}, \mathbf{\Delta}^{-1}\mathbf{\Lambda}\mathbf{\Delta}^{-1}), \end{aligned} \quad (6)$$

with  $\Gamma(u_0)$  given by (4),

$$\mathbf{\Delta} = f_U(u_0) \begin{pmatrix} 1 & \mu_1 \\ \mu_1 & \mu_2 \end{pmatrix} \otimes \Gamma(u_0) \quad (7)$$

and

$$\mathbf{\Lambda} = f_U(u_0) \begin{pmatrix} \nu_0 & \nu_1 \\ \nu_1 & \nu_2 \end{pmatrix} \otimes \Gamma(u_0). \quad (8)$$

Furthermore, if  $K(\cdot)$  is symmetric, then

$$\sqrt{nh} \left[ \hat{\mathbf{a}}_{MLE}(u_0) - \mathbf{a}(u_0) - \frac{h^2\mu_2}{2} \mathbf{a}''(u_0) + o_p(h^2) \right] \xrightarrow{\mathcal{D}} \mathbf{N}(\mathbf{0}, \mathbf{\Sigma}(u_0)), \quad (9)$$

where

$$\mathbf{\Sigma}(u_0) = \nu_0\mathbf{\Gamma}^{-1}(u_0)/f_U(u_0). \quad (10)$$

Note that the bias and variance expressions in Theorem 1 can be deduced from the general theorem from Carroll et al. (1998). However, the main difference is that here we establish the results in terms of asymptotic normality, whereas those authors established them for the general case using conditional expectations.

*Theorem 2.* Under the assumptions in Theorem 1,  $\hat{\beta}_{OS}$  has the same asymptotic distribution as  $\hat{\beta}_{MLE}$ , provided that the initial estimator  $\hat{\beta}_0$  satisfies

$$\mathbf{H}(\hat{\beta}_0 - \beta) = O_p\{h^2 + (nh)^{-1/2}\}. \quad (11)$$

Consequently, the fully iterative MLE and the one-step estimate share the same asymptotic properties provided that (11) is fulfilled, which provide the theoretical basis for using the one-step approach in practice. The asymptotic MSE of the two estimators  $\hat{a}_{j,MLE}(u_0)$  and  $\hat{a}_{j,OS}(u_0)$  is

$$\text{MSE} = \frac{h^4}{4} \mu_2^2 \{a''_j(u_0)\}^2 + \frac{\nu_0\sigma_{jj}^2(u_0)}{nhf_U(u_0)}$$

when  $K(\cdot)$  is symmetric, where  $\sigma_{jj}^2(u_0)$  is the  $j$ th diagonal element of  $\mathbf{\Gamma}^{-1}(u_0)$ . Then the MSE is of order  $n^{-4/5}$  if the optimal bandwidth  $[\nu_0\sigma_{jj}^2(u_0)/\{\mu_2^2(a''_j(u_0))^2 f_U(u_0)\}]^{1/5} n^{-1/5}$  is used.

### 2.4 Standard Errors

Because the local likelihood (2) is a weighted likelihood function of a parametric generalized linear model, the covariance matrix of  $\hat{\beta}_{MLE}$  can be estimated from conventional techniques. Let  $q_j(s, y) = (\partial^j/\partial s^j)l\{g^{-1}(s), y\}$  and

$$\begin{aligned} \hat{\Gamma}(u_0) \\ = -\frac{1}{n} \sum_{i=1}^n q_2 \left[ \sum_{j=1}^p \{\hat{a}_j(u_0)X_{ij} + \hat{b}_j(u_0)(U_i - u_0)\}, Y_i \right] \\ \times K_h(U_i - u_0) \left( \frac{\mathbf{X}_i}{\mathbf{X}_i(U_i - u_0)/h} \right)^{\otimes 2}, \end{aligned} \quad (12)$$

Table 1. Bivariate Summary of Simulation Results for the Logistic Regression Model

n	h	MLE		One-step			Two-step		
		$\mu$	$\sigma$	$\mu$	$\sigma$	$\rho_*$	$\mu$	$\sigma$	$\rho_*$
400	.10	2.2278	2.0874	1.8537	.9759	.8656	2.1244	1.5315	.8274
	.20	1.0669	.4491	1.0576	.4378	.9991	1.0669	.4491	1.0000
	.40	.9454	.1600	.9447	.1593	1.0000	.9454	.1600	1.0000
	.075	1.2451	.6639	1.1644	.3767	.8342	1.2256	.5301	.9656
800	.15	.7280	.2573	.7234	.2459	.9993	.7280	.2573	1.0000
	.30	.7433	.1009	.7429	.1005	1.0000	.7433	.1009	1.0000



Table 2. Standard Deviations of Estimators for Logistic Regression Model

n	h	u	$\hat{a}_0(u)$		$\hat{a}(u)$		$\hat{a}_2(u)$	
			SD	SD <sub>a</sub> (SD <sub>std</sub> )	SD	SD <sub>a</sub> (SD <sub>std</sub> )	SD	SD <sub>a</sub> (SD <sub>std</sub> )
400	.2	.25	.3185	.2673 (.0470)	.4890	.4069 (.0776)	.5082	.3986 (.0893)
		.50	.3410	.2782 (.0451)	.5413	.4330 (.0809)	.4135	.3568 (.0591)
		.75	.4315	.3542 (.0776)	.5372	.4542 (.0996)	.5809	.4431 (.0969)
400	.3	.25	.2294	.2051 (.0231)	.3424	.3201 (.0447)	.3317	.2956 (.0403)
		.50	.2570	.2315 (.0315)	.3931	.3538 (.0527)	.3490	.3122 (.0431)
		.75	.2850	.2686 (.0423)	.3929	.3581 (.0557)	.3788	.3328 (.0500)
800	.15	.25	.2418	.2214 (.0214)	.3638	.3460 (.0501)	.3804	.3486 (.0532)
		.50	.2249	.2196 (.0233)	.4040	.3569 (.0512)	.3124	.2812 (.0356)
		.75	.3146	.2928 (.0478)	.4209	.3804 (.0667)	.3987	.3781 (.0631)

where  $\mathbf{A}^{\otimes 2}$  denotes  $\mathbf{A}\mathbf{A}^T$  for a matrix or vector  $\mathbf{A}$ . Then the covariance matrix of  $\hat{\beta}_{MLE}$  can be estimated as

$$\hat{\Sigma}^*(u_0) = \hat{\Gamma}(u_0)^{-1} \hat{\Lambda}(u_0) \hat{\Gamma}(u_0)^{-1}, \tag{13}$$

where

$$\begin{aligned} \hat{\Lambda}(u_0) &= \frac{h}{n} \sum_{i=1}^n q_1^2 \left[ \sum_{j=1}^p \{ \hat{a}_j(u_0) X_{ij} + \hat{b}_j(u_0) (U_i - u_0) \}, Y_i \right] \\ &\quad \times K_h^2(U_i - u_0) \left( \begin{matrix} \mathbf{X}_i \\ \mathbf{X}_i(U_i - u_0)/h \end{matrix} \right)^{\otimes 2}. \end{aligned}$$

In the implementation in Section 3, a ridge regression technique is used, and hence the matrix  $\hat{\Gamma}(u_0)$  in (13) is modified slightly to reflect this change.

The explicit formula for the asymptotic covariance matrix in (10) provides an alternative estimate of the asymptotic covariance matrix of  $\mathbf{a}(u_0)$  (not the full vector  $\hat{\beta}_{MLE}$ )  $\Sigma(u_0)$ . Therefore, a direct estimate of  $\Sigma(u_0)$  is  $\tilde{\Sigma}(u_0) = \nu_0 \hat{\Gamma}_S(u_0)^{-1}$ , where  $\hat{\Gamma}_S(u_0)$  is the  $p \times p$  upper corner submatrix of  $\hat{\Gamma}(u_0)$  given by (12).

### 2.5 Hypothesis Testing

When fitting a varying-coefficient model, one naturally asks whether the coefficient functions are actually varying or whether any particular covariate is significant in the model. For simplicity of description, we consider only the

first hypothesis testing problem

$$H_0: a_1(u) \equiv a_1, \dots, a_p(u) \equiv a_p, \tag{14}$$

though the technique also applies to other testing problems. A useful procedure is based on the nonparametric likelihood ratio test statistic

$$T = 2\{l(H_1) - l(H_0)\}, \tag{15}$$

where  $l(H_0)$  and  $l(H_1)$  are the log-likelihood functions computed under the null and the alternative hypotheses.

For parametric models, the likelihood ratio statistic follows asymptotically a chi-squared-distribution with degrees of freedom  $f - r$ , where  $r$  and  $f$  are the number of parameters under the null and alternative hypotheses. For the nonparametric alternative, the effective number of parameters  $f$  tends to infinite. Thus the test statistic will be asymptotically normal, independent of the values  $a_1, \dots, a_p$ . For the rigorous justification, we refer to the article by Fan, Zhang, and Zhang (1999) that considered sieve likelihood ratio tests in a general setting and demonstrated that the Wilks type of phenomenon holds for a large variety of nonparametric problems. This in turn suggests that we can use the following conditional bootstrap to construct the null distribution of  $T$ . Let  $\{\hat{a}_j\}$  be the MLE under the null hypothesis. Given the covariates  $(U_i, \mathbf{X}_i)$ , generate a bootstrap sample  $Y_i^*$  from the given distribution of  $Y$  with the estimated linear predictor  $\hat{\eta}(U_i, \mathbf{X}_i) = \sum_{j=1}^p \hat{a}_j X_{ij}$ , and compute the test statistic  $T^*$  in (15). Use the distribution of  $T^*$  as an approximation to the distribution of  $T$ . This method is valid because the asymptotic null distribution does not depend on the values of  $\{a_j\}$  (Fan et al. 1999).

Note that the foregoing conditional bootstrap method applies readily to the Poisson and Bernoulli distributions, because in these cases the distribution of  $Y$  does not involve any dispersion parameters. It is really a simulation approximation to the conditional distribution of  $T$  given observed covariates under the particular null hypothesis,  $H_0: a_j(u) = \hat{a}_j$  ( $j = 1, \dots, p$ ). As pointed out earlier, this approximation is valid under both  $H_0$  and  $H_1$ , as the null distribution does not asymptotically depend on the values of  $\{a_j\}$ . In the case where model (1) involves a dispersion parameter (e.g., the Gaussian model), the dispersion parameter should be estimated based on the residuals from the alternative hypothesis. This is again due to the Wilks type of results demonstrated by Fan et al. (1999).

Table 3. Six Empirical Percentiles for the Logistic Model

	10	25	50	75	90	95
<b>Conditional bootstrap</b>						
	7.9579	10.7189	14.2569	18.2625	22.2566	24.9903
	8.2450	11.0170	14.6601	18.4897	22.4177	25.5829
	8.0004	10.9871	14.2667	18.0413	22.5517	25.1661
	8.7738	11.4311	14.8061	18.5209	22.7029	25.3781
	8.7906	11.4672	14.9130	18.6168	22.3256	24.7104
<b>Unconditional bootstrap</b>						
	7.6381	10.7167	14.5487	18.6276	22.2205	24.4597
	7.3478	10.1290	13.9934	17.9622	21.8270	24.4429
	7.7238	11.3849	14.6151	18.4796	22.5899	24.7270
	8.8042	11.3762	14.8076	18.7571	22.0560	25.1550
	8.7865	11.3472	14.5975	18.5198	23.1476	25.8297




Figure 2. Simulation Results for Example 2 With Sample Size 200. The caption is the same as for Figure 1.

For testing the hypothesis such as  $a_p(\cdot) = 0$ , the foregoing conditional bootstrap idea continues to apply. In this case, the data should be generated from the mean function  $g\{m(\mathbf{u}, \mathbf{x})\} = \sum_{j=1}^{p-1} \hat{a}_j(\mathbf{u})x_j$ , where  $\hat{a}_j(\cdot)$  is an estimate under the alternative hypothesis.

### 2.6 Implementation of One-Step Local Maximum Likelihood Estimation

Suppose that we wish to evaluate the functions  $\hat{a}(\cdot)$  at grid points  $u_j, j = 1, \dots, n_{\text{grid}}$ . Our idea for finding initial estimators is as follows. Take a point  $u_{i_0}$ , usually the cen-

ter of the grid points. Compute the local MLE,  $\hat{\beta}_{\text{MLE}}(u_{i_0})$ . Use this estimate as the initial estimate for the point  $u_{i_0+1}$ , and apply (3) to obtain  $\hat{\beta}_{\text{OS}}(u_{i_0+1})$ . Now use  $\hat{\beta}_{\text{OS}}(u_{i_0+1})$  as the initial estimate at the point  $u_{i_0+2}$ , and apply (3) to obtain  $\hat{\beta}_{\text{OS}}(u_{i_0+2})$  and so on. Likewise, we can compute  $\hat{\beta}_{\text{OS}}(u_{i_0-1}), \hat{\beta}_{\text{OS}}(u_{i_0-2})$ , and so on. In this way, we obtain our estimates at all grid points.

There are a couple of possible variations to the foregoing technique. The first variation is to calculate a fresh local MLE as a new initial value after iterating along the grid points for a while; for example, if we wish to evaluate the

Table 4. Bivariate Summary of Simulation Output for the Poisson Regression Model

n	h	MLE		One-step			Two-step		
		$\mu$	$\sigma$	$\mu$	$\sigma$	$\rho_*$	$\mu$	$\sigma$	$\rho_*$
200	.075	.3632	.0692	.3468	.0562	.8691	.3632	.0692	1.0000
	.15	.3220	.0510	.3202	.0504	.9925	.3220	.0510	1.0000
	.30	.5852	.0425	.5835	.0426	.9990	.5852	.0425	1.0000
400	.075	.2309	.0352	.2279	.0347	.9866	.2309	.0352	1.0000
	.15	.2581	.0325	.2571	.0322	.9942	.2581	.0325	1.0000
	.30	.5603	.0292	.5581	.0293	.9988	.5603	.0292	1.0000

functions at 200 grid points and are willing to compute the local maximum likelihood at five distinct points. A sensible placement of these points is  $u_{20}, u_{60}, u_{100}, u_{140}$ , and  $u_{180}$ . Use, for example,  $\hat{\beta}_{MLE}(u_{60})$ , along with the idea in the last paragraph, to compute  $\hat{\beta}_{OS}(u_i)$  for  $i = 40, \dots, 79$ . In our implementation, this modified technique is used.

Another useful modification is to use a two-step method. We use the scenarios given in the preceding paragraph as an illustration. After obtaining  $\hat{\beta}_{MLE}(u_{60})$ , say, we apply (3) to obtain  $\hat{\beta}_{OS}(u_{61})$ . Regarding  $\hat{\beta}_{OS}(u_{61})$  as an initial value, we use (3) to obtain a “two-step” estimator  $\hat{\beta}_{TS}(u_{61})$ . We then use  $\hat{\beta}_{TS}(u_{61})$  as an initial value for the grid point  $u_{62}$  and iterate (3) twice to obtain  $\hat{\beta}_{TS}(u_{62})$ , and so on. This implementation requires approximately twice as much effort to compute the estimates as the one-step method; however, our empirical studies show that there are no significant differences between the two procedures. Section 3 provides details.

The theoretical basis for the foregoing “one-step” and “two-step” procedures is as follows. When the grid points are sufficiently fine,  $\hat{\beta}_{MLE}(u_{i_0})$  will be very close to  $\hat{\beta}_{MLE}(u_{i_0+1})$ . Indeed, when the grid span is of order  $O\{h_n^2 + (nh_n)^{-1/2}\}$  which usually is true for most applications,  $\hat{\beta}_{MLE}(u_{i_0})$  satisfies the condition given in Theorem 2. Therefore,  $\hat{\beta}_{OS}(u_{i_0+1})$  is as efficient as the fully iterative local MLE at the point  $u_{i_0+1}$ . Using the same reasoning,  $\hat{\beta}_{OS}(u_{i_0+2})$  is as efficient as the local MLE at the point  $u = u_{i_0+2}$ , and so on. The same arguments are still applicable for the two-step estimator. A fresh start is needed because of stochastic error accumulation as iterations along grid points march on.

Based on the foregoing theoretical considerations, we suggest a very simple rule of thumb for choosing the number of grid points:  $n_{grid} = \max\{200, IQR^2/h^2\}$ , where IQR is the interquantile range of  $U_1, \dots, U_n$ . In such a way, ap-

proximation errors between estimates at two consecutive grid points are of order  $O(h^2)$ , satisfying the critical condition (11).

### 3. SIMULATIONS AND APPLICATIONS

In this section we first discuss how to implement the one-step procedure for the Bernoulli and Poisson models. We then illustrate the performance of the proposed one-step method and compare it to the two-step estimator and the fully iterative local MLE. The performance of estimator  $\hat{a}(\cdot)$  is assessed via the square root of average squared errors (RASE),

$$RASE^2 = n_{grid}^{-1} \sum_{j=1}^p \sum_{k=1}^{n_{grid}} \{\hat{a}_j(u_k) - a_j(u_k)\}^2, \quad (16)$$

where  $\{u_k, k = 1, \dots, n_{grid}\}$  are the grid points at which the functions  $\{a_j(\cdot)\}$  are estimated.

In the following two simulated examples, the covariates  $X_1$  and  $X_2$  are standard normal random variables with correlation coefficient  $2^{-1/2}$ , and  $U$  is uniformly distributed over  $[0, 1]$ , independent of  $(X_1, X_2)$ . Three bandwidths are used to represent widely varying degrees of smoothness. Over this range of bandwidths, we compare the performances among the one-step, the two-step, and the fully iterative local MLE methods. We use the Epanechnikov kernel  $K(u) = .75(1 - u^2)_+$  and  $n_{grid} = 200$ .

#### 3.1 Logistic Regression

For a Bernoulli distribution, the one-step estimator is given by

$$\hat{\beta}_{OS} = \hat{\beta}_0 + \begin{pmatrix} \mathbf{H}_{n,0} & \mathbf{H}_{n,1} \\ \mathbf{H}_{n,1} & \mathbf{H}_{n,2} \end{pmatrix}^{-1} \begin{pmatrix} \mathbf{v}_{n,0} \\ \mathbf{v}_{n,1} \end{pmatrix}, \quad (17)$$

where  $\mathbf{H}_{n,j} = \sum_{i=1}^n K_h(U_i - u_0) \hat{p}_{i0} (1 - \hat{p}_{i0}) (U_i - u_0)^j \mathbf{X}_i \mathbf{X}_i^T, j = 0, 1, 2, \hat{p}_{i0}$  satisfies logit  $(\hat{p}_{i0}) = \sum_{j=1}^p \{\hat{a}_{j,0} + \hat{b}_{j,0}(U_i - u_0)\} X_{ij}$ , and  $\mathbf{v}_{n,j} = \sum_{i=1}^n K_h(U_i - u_0) (Y_i - \hat{p}_{i0})$

Table 5. Standard Deviations of Estimators for the Poisson Regression Model

n	h	u	$\hat{a}_0(u)$		$\hat{a}_1(u)$		$\hat{a}_2(u)$	
			SD	$SD_a (SD_{std})$	SD	$SD_a (SD_{std})$	SD	$SD_a (SD_{std})$
200	.15	.25	.0105	.0092 (.0013)	.0148	.0118 (.0024)	.0156	.0126 (.0026)
		.50	.0094	.0088 (.0011)	.0148	.0112 (.0022)	.0150	.0118 (.0024)
		.75	.0100	.0088 (.0011)	.0142	.0112 (.0023)	.0151	.0119 (.0023)
400	.075	.25	.0094	.0085 (.0012)	.0130	.0106 (.0021)	.0136	.0107 (.0022)
		.50	.0093	.0083 (.0011)	.0127	.0104 (.0022)	.0130	.0105 (.0021)
		.75	.0090	.0081 (.0011)	.0137	.0101 (.0022)	.0133	.0102 (.0022)

Table 6. Six Empirical Percentiles for the Poisson Model

10	25	50	75	90	95
<b>Conditional bootstrap</b>					
12.1646	15.1401	18.6981	22.6260	26.1432	28.8494
11.7506	14.5010	18.0994	22.3809	26.1237	29.4936
11.7946	14.7005	18.3495	22.2918	26.0064	29.2165
11.4662	14.6917	18.2475	22.4623	27.0587	29.6887
11.9894	14.7869	18.5571	22.3593	26.7014	29.7923
<b>Unconditional bootstrap</b>					
11.9492	14.7920	18.5509	22.3383	26.7474	28.8094
11.1599	14.7156	18.7054	22.2915	26.6170	28.9831
11.4378	14.8132	18.4080	22.3890	26.5858	29.4816
11.8238	14.6817	18.5090	22.7050	26.4776	29.3814
11.8365	14.9721	18.7674	22.9402	26.5929	28.9815

$(U_i - u_0)^j \mathbf{X}_i, j = 0, 1$ . The two-step estimator  $\hat{\beta}_{TS}$  is obtained by iterating (17) twice, and the local MLE is obtained by iterating (17) until convergence.

In practice, the matrix in (17) can be singular or nearly singular when the local data are sparse. To attenuate this difficulty, one may follow the idea of ridge regression (Fan and Chen 1999; Seifert and Gasser 1996). Then an issue arises as to how to choose the ridge parameters. Note that the  $k$ th diagonal element of  $\mathbf{H}_{n,j}$  ( $j = 0$  and  $2$ ) is approximately of order

$$E(X_k^2|U = u_0)\hat{p}_0(1 - \hat{p}_0)h^{j-1} \int u^j K(u) du N$$

with

$$\hat{p}_0 = \frac{\exp(\hat{\mathbf{a}}_0^T \bar{\mathbf{X}})}{1 + \exp(\hat{\mathbf{a}}_0^T \bar{\mathbf{X}})}, \tag{18}$$

where  $N = nh f_U(u_0)$  and  $\bar{\mathbf{X}} = \frac{1}{n} \sum_{i=1}^n \mathbf{X}_i$ . The parameter  $N$  can be intuitively understood as the effective number of local data points. This motivates us to use the ridge parameter

$$r_{j,k} = \left( \frac{1}{n} \sum_{i=1}^n X_{ik}^2 \right) \hat{p}_0(1 - \hat{p}_0)h^{j-1} \int u^j K(u) du$$

for the  $k$ th diagonal element of  $\mathbf{H}_{n,j}$ . Using such a ridge parameter will not alter the asymptotic behavior and will prevent the matrix from becoming nearly singular when  $N$  is small. However, it affects, and indeed ameliorates, the finite-sample properties of estimators for small sample sizes.

*Example 1.* Take  $\mathbf{X} = (1, X_1, X_2)^T$ , and the coefficient functions in (1) are given by

$$a_0(u) = \exp(2u - 1), \quad a_1(u) = 8u(1 - u), \tag{19}$$

and


$$a_2(u) = 2 \sin^2(2\pi u). \tag{20}$$

Figure 1(a) depicts the marginal distributions for the ratios of the overall RASE defined in (16), using three bandwidths  $h = .1, .2$ , and  $.4$ . It is evident that the performance of the


one-step, two-step, and fully iterative estimators are comparable for a wide range of bandwidths. As expected, the performance of the two-step estimator is closer to that of the local MLE. Figures 1(b)–(d) give the estimate of the coefficient functions from a typical sample. The typical sample is selected in such a way that its RASE value is the median in the 400 RASE values. Table 1 summarizes the simulation results with  $\mu$  and  $\sigma$  denoting the mean and standard deviation of the RASE in 400 simulations. Here  $\rho_*$  indicates the correlation coefficient between the RASE of the MLE and the RASE of the one-step (or two-step) method. Note that the correlation coefficients are close to 1, which indicates that the one-step and two-step methods closely follow the MLE. Note also that the larger the bandwidths, the larger the correlation coefficients. This is due to the fact that a larger bandwidth implies more local data points, which makes the asymptotic theory more relevant. As expected, the correlation coefficients for the two-step method are larger than those of the one-step method, because the former is closer to the MLE.

We now test the accuracy of our standard error formula (13). The standard deviation, denoted by SD in Table 2, of 400 estimated  $\hat{a}_j(u_0)$ , based on 400 simulations, can be regarded as the true standard errors. The average and the standard deviation of 400 estimated standard errors, denoted by  $SD_a$  and  $SD_{std}$ , summarize the overall performance of the standard error formula (13). Table 2 presents the results at the points  $u_0 = .25, .50$ , and  $.75$ . It suggests that our standard error formula somewhat underestimates the true standard deviation, though the difference is within two standard deviations of the Monte Carlo errors. The bias becomes smaller as the number of local data points  $nh_n$  increases (see the last two situations). This is consistent with our asymptotic theory.


Next, we conduct a simulation study to see whether the asymptotic null distribution of the test statistic  $T$  defined in (15) depends on the values of  $\{a_j\}$  under  $H_0$  [see (14)] and the limiting conditional null distributions are dependent on the covariate values. To this end, we compute the unconditional null distribution of  $T$  with  $n = 400$ , via 1,000 Monte Carlo simulations, for five different sets of values of  $\{a_j\}$ . These sets of parameters are quite far apart. The resulting five densities are depicted in Figure 1(e) (thick curves). They are very close, suggesting that the asymptotic null distribution is not very sensitive to the values of  $\{a_j\}$ . To validate our conditional bootstrap method, five typical datasets were selected from our previous 400 simulations. The estimated conditional bootstrap null distributions, based on 1,000 bootstrap samples, are plotted as thin curves in Figure 1(e). Six empirical percentiles for five different sets of values of  $\{a_j\}$  and covariates are listed in Table 3. Both Figure 1(e) and Table 3 show that they are very close to the true null distribution. This demonstrates empirically that our bootstrap method gives a reasonably good approximation to the true null distribution even when the data were generated from an alternative model (19)–(20).



(a)



(b)



(c)

Figure 3. (a) Scatterplot of Log Transformation of the Environmental Dataset Studied in Section 3.3, Where the Curve is the Estimate of  $a_1(t) + a_2(t)\bar{x}_1 + a_3(t)\bar{x}_2 + a_4(t)\bar{x}_3$ , Where  $\bar{x}_i$  is the Average Pollutant Level  $x_i$ ; (b) Plot of the Cross-Validation Functions  $CV_1(h)$  (Solid Line) and  $CV_2(h)$  (Dashed and Dotted Line) Against the Bandwidth; (c) The Same as in (b), but With the Cross-Validation Based on Random Partitions of the Dataset.






Figure 4. The Estimated Coefficient Functions via the One-Step Approach With Bandwidth Chosen by the CV. The dashed curves are the estimated function plus/minus twice estimated standard errors. (a) Coefficient function  $a_1(t)$ ; (b) coefficient function  $a_2(t)$ ; (c) coefficient function  $a_3(t)$ ; (d) coefficient function  $a_4(t)$ .

To examine the power of the proposed test, we consider the null hypothesis

$$H_0: a_j(u) = \theta_j, \quad j = 0, 1, 2,$$

versus

$$H_1: a_j(u) \neq \theta_j, \quad \text{for at least one } j.$$

The power functions are evaluated under a sequence of the alternative models indexed by  $\beta$ ,

$$H_1: a_j(u) = a_{j0} + \beta(a_j^0(u) - a_{j0}), \quad j = 0, 1, 2, \quad (0 \leq \beta \leq .8),$$

where  $\{a_j^0(u)\}$  are given in (19)–(20) and  $a_{j0} = E\{a_j(U)\}$ . Figure 1(f) depicts the five power functions based on 1,000 simulations for the sample size  $n = 400$  at five different significance levels: .5, .25, .10, .05, and .01. When  $\beta = 0$ , the special alternative collapses into the null hypothesis. The powers at  $\beta = 0$  for the foregoing five significance levels are .532, .281, .101, .047, and .012. This shows that the conditional bootstrap method gives the right levels of test. The power functions increase rapidly as  $\beta$  increases. This in turn shows that the test proposed in Section 2.5 works well.

### 3.2 Poisson Regression

For a Poisson model with the canonical link, by straightforward calculation, the one-step estimator is given similarly to (17) but now  $\mathbf{H}_{n,j} = \sum_{i=1}^n K_h(U_i - u_0) \hat{\lambda}_{i0} (U_i - u_0)^j \mathbf{X}_i \mathbf{X}_i^T$ ,  $j = 0, 1, 2$ ,  $\hat{\lambda}_{i0} = \exp\{\sum_{j=1}^p \{\hat{a}_{j0} + \hat{b}_{j0}(U_i - u_0)\} X_{ij}\}$ , and  $\mathbf{v}_{n,j} = \sum_{i=1}^n K_h(U_i - u_0) (Y_i - \hat{\lambda}_{i0}) (U_i - u_0)^j \mathbf{X}_i$ ,  $j = 0, 1$ . Using the same arguments as in the previous

section, the ridge parameters

$$r_{j,k} = \left( \frac{1}{n} \sum_{i=1}^n X_{ik}^2 \right) \hat{\lambda}_0 h^{j-1} \int u^j K(u) du$$

with

$$\hat{\lambda}_0 = \exp(\hat{\mathbf{a}}_0^T \bar{\mathbf{X}}) \tag{21}$$

are used to alleviate the possible singularity of matrix  $\mathbf{H}_{n,j}$  ( $j = 0$  and  $2$ ) in (17).

*Example 2.* The conditional distribution of  $Y$  given covariates  $U$ ,  $X_1$ , and  $X_2$  is taken to be Poisson with the linear predictor

$$\eta(u, \mathbf{x}) = 5.5 + .1\{a_0(u) + a_1(u)x_1 + a_2(u)x_2\},$$

where the coefficient functions  $a_0(u)$ ,  $a_1(u)$ , and  $a_2(u)$  are the same as those in Example 1. The coefficients 5.5 and .1 are chosen so that the range of simulated data is close to that of the environmental data in Section 3.3.

Figure 2 and Table 4 summarize the result for  $n = 200$ . It shows again that the one-step, two-step, and iterative local MLE have comparable performance. A typical estimated function with bandwidth  $h = .15$  is presented in Figures 2(b)–(d). Because of different signal-to-noise ratios, the functions here are indeed estimated better than those given in Example 1. Similar to Example 1, the performance of our estimated standard error formula (13) is summarized in Table 5. Clearly, our estimated standard errors are very close to the true ones.

Similar to Example 1, the procedure of testing hypothesis is applied to this example. Both unconditional and conditional estimated densities of  $T$  are displayed in Figure 2(e). Six empirical percentiles are listed in Table 6. The corresponding power functions are presented in Figure 2(f). The




Figure 5. (a) Time Series Plot of Pearson's Residuals and (b) Plot of the Autocorrelation Coefficients Versus Time Lag. The two dashed curves are  $\pm 1.96/\sqrt{n}$ , where  $n$  is the sample size.

same conclusions as those for Example 1 can be drawn for the Poisson regression model. In particular, the test has the correct levels of significance; see the power functions in Figure 2(e) at  $\beta = 0$ .

### 3.3 Real Data Examples

*Example 3.* We illustrate in this example our proposed procedure via an application to the environmental dataset mentioned in Section 1. Of interest is in studying the association between levels of pollutants and the number of total hospital admissions for circulatory and respiratory problems on every Friday from January 1, 1994, to December 31, 1995 and to examine the extent to which the associa-

tion varies over time. The covariates are taken as the levels of pollutants sulfur dioxide,  $X_2$  (in  $\mu\text{g}/\text{m}^3$ ), nitrogen dioxide,  $X_3$  (in  $\mu\text{g}/\text{m}^3$ ) and dust,  $X_4$  (in  $\mu\text{g}/\text{m}^3$ ). Because the admissions "events" occur at certain points in time, it is reasonable to model the number of admissions as a Poisson process and use the Poisson regression model with the mean  $\lambda(t, \mathbf{x})$  given by

$$\log\{\lambda(t, \mathbf{x})\} = a_1(t) + a_2(t)x_2 + a_3(t)x_3 + a_4(t)x_4. \quad (22)$$

A multifold cross-validation method is used to select a bandwidth. We partition the data into  $Q$  groups, with the  $j$ th group consisting of data points with indices

$$d_j = \{20k + j, k = 1, 2, \dots\}, \quad j = 0, \dots, Q - 1.$$

For each  $j$ , the  $j$ th group of data is deleted, and model (22) is fitted for the remaining data. Then the deviance (McCullagh and Nelder 1989, p. 34) or the sum of squares of Pearson's residuals is computed. This leads to two cross-validation criteria,

$$CV_1(h) = \sum_{j=0}^{Q-1} \sum_{i \in d_j} 2[y_i \log\{y_i/\hat{y}_{-d_j}(U_i, \mathbf{X}_i)\} - \{y_i - \hat{y}_{-d_j}(U_i, \mathbf{X}_i)\}]$$

and

$$CV_2(h) = \sum_{j=0}^{Q-1} \sum_{i \in d_j} \left\{ \frac{y_i - \hat{y}_{-d_j}(U_i, \mathbf{X}_i)}{\sqrt{\hat{y}_{-d_j}(U_i, \mathbf{X}_i)}} \right\}^2,$$

where  $\hat{y}_{-d_j}(U_i, \mathbf{X}_i)$  is a fitted value with the data in  $d_j$  deleted. In the implementation, we choose  $Q = 20$ . Figure 3(b) depicts the cross-validation functions  $CV_1(h)$  and  $CV_2(h)$ , which give the optimal bandwidth  $h = .1440 \times 105$ . To see how sensitive the aforementioned partition is to the CV curves, the dataset is randomly partitioned into 20 groups, and then CV scores are computed based on the procedure described earlier. The results are depicted in Figure 3(c), which, in conjunction with Figure 3(b), shows that the CV functions is not very sensitive to the partition. The estimated coefficient functions based on the one-step procedure are summarized in Figure 4, because the results based on both the one-step and the fully iterative methods are very close. They describe the extent to which the association between the pollutants and the number of hospital admissions vary over time. The figure shows clearly that the coefficient functions vary with time. The two dashed curves are the estimated function plus/minus twice the estimated standard errors. They give us an idea of the pointwise confidence intervals with bias ignored.

A question arises as to whether or not the data are highly correlated. To check for the serial correlation, Pearson's residuals are computed. The time series plot of the residuals is given in Figure 5(a), and the plot of the corresponding autocorrelation coefficients against time lag is presented in Figure 5(b). There is no pattern in Figure 5(a); thus Figure 5(a) together with Figure 5(b) leads to the conclusion that there is no evidence that the data are serially correlated.

We now apply the procedure proposed in Section 2.5 to testing whether the coefficients are actually time varying. The MLE under the null hypothesis is (5.4499, -.0025, .0015, -.0005) with estimated standard deviation (.0195, .0006, .0006, .0005). The test statistic (15) is  $T = 389.41$ . Based on 1,000 bootstrap replications, the sample mean and sample variance of  $T^*$  are 26.64 and 48.40. The distribution of  $T$  is approximated by a chi-square distribution with degrees of freedom 27 (see Fig. 6). The  $p$  value is close to 0, which strongly rejects the null hypothesis. Therefore, it suggests that the varying-coefficient model gives a much better fit than the parametric model.

Now we use our testing approach proposed in Section 2.5 to check whether there is any covariate that can be deleted from the model. We start with  $X_4$ , because the parametric Poisson model concludes that the dust level ( $X_4$ ) is not statistically significant. To examine whether the variable  $X_4$  is significant in the varying-coefficient model, we apply the idea in Section 2.5 to testing the hypothesis; the function  $a_4(\cdot)$  is 0. The maximum likelihood ratio test statistic is  $T = 20.1847$ . Based on 1,000 bootstrap samples, the  $p$  value is .321 (and the sample mean and variance of  $T^*$  are 17.7352 and 37.1976). Therefore, the variable  $X_4$  can be dropped from the varying-coefficient model. After deleting the variable dust level ( $X_4$ ), we apply the same procedure to test whether  $X_3$  is statistically significant in the varying-coefficient model; that is, to test  $H_0$ :




Figure 6. The Estimated Density of  $T$  by Monte Carlo Simulation. The solid curve is the estimated density, and the dashed curve represents the density of chi-squared distribution with degrees of freedom 27.




Figure 7. Estimated Coefficient Functions (Solid Curves) via the One-Step Approach With Bandwidth Chosen by the CV. The dot curves are the estimated functions plus/minus twice estimated standard errors. (a) Coefficient of  $a_1(\cdot)$ ; (b) estimated coefficient of  $a_2(\cdot)$ ; (c) estimated coefficient of  $a_3(\cdot)$ ; (d) estimated coefficient of  $a_4(\cdot)$ .

$\log\{\lambda(t, \mathbf{x})\} = a_1(t) + a_2(t)x_2$  against  $H_1: \log\{\lambda(t, \mathbf{x})\} = a_1(t) + a_2(t)x_2 + a_3(t)x_3$ . As a result, the maximum likelihood ratio test statistic is  $T = 39.7473$ , and the  $p$  value is .039 (with sample mean and variance of  $T^*$  of 27.5071 and 39.5808), based on 1,000 bootstrap samples. Therefore, the variable nitrogen dioxide ( $X_3$ ) is significant at the significant level .05. By the same token, the variable sulfur dioxide ( $X_2$ ) is significant as well.

*Example 4.* Now we apply the methodology proposed in this article to analyze the dataset, burns data collected by General Hospital Burn Center at the University of Southern California. The binary response variable  $Y$  is 1 for those victims who survived their burns and 0 otherwise, and covariates  $X_1 = \text{age}$ ,  $X_2 = \text{sex}$ ,  $X_3 = \log(\text{burn area} + 1)$ , and binary variable  $X_4 = \text{oxygen}$  (0 if oxygen supply is normal, 1 otherwise) are considered. We are interested in studying how burn areas and the other variables affect survival probabilities for victims in different age groups. This naturally leads to the varying-coefficient model

$$\text{logit}\{p(x_1, x_2, x_3, x_4)\} = a_1(x_1) + a_2(x_1)x_2 + a_3(x_1)x_3 + a_4(x_1)x_4. \quad (23)$$

Figure 7 presents the estimated coefficients for model (23) via the one-step approach with bandwidth  $h = 65.7882$ , selected by a CV method.

A natural question arises whether the coefficients in (23) are actually varying. To see this, we consider the parametric logistic regression model

$$\text{logit}\{p(x_1, x_2, x_3, x_4)\} = \beta_0 + \beta_1x_1 + \beta_2x_2 + \beta_3x_3 + \beta_4x_4 \quad (24)$$

as the null model. As a result, the MLE of  $(\beta_0, \dots, \beta_4)$  in model (24) and its standard deviation are (23.2213, -6.1485, -.4661, -2.4496, -.9683) and (1.9180, .6647, .2825, .2206,

.2900). The test statistic  $T$  proposed in Section 2.5 is 54.9601 with a  $p$  value of 0, based on 1,000 bootstrap samples (with sample mean and variance of  $T^*$  of 5.9756 and 10.7098). This implies that the varying-coefficient logistic regression model fits the data much better than the parametric fit. It also allows us to examine the extent to which the regression coefficients vary over different ages.

To examine whether there is any gender gap for different age groups or whether the variable  $X_4$  affects the survival probabilities for burn victims of different ages, we consider testing hypothesis  $H_0$ : both  $a_2(\cdot)$  and  $a_4(\cdot)$  are constant under model (23). The corresponding test statistic  $T$  is 3.2683 with a  $p$  value of .7050, based on 1,000 bootstrap samples. This in turn suggests that the coefficient functions  $a_2(\cdot)$  and  $a_4(\cdot)$  are independent of age and indicates that there are no gender differences for different age groups.

Finally, we examine whether the covariates sex and oxygen are statistically significant in model (23). The likelihood ratio test for this problem is  $T = 11.2727$  with a  $p$  value of .0860, based on 1,000 bootstrap samples (with sample mean and variance of  $T^*$  of 5.2867 and 9.7630). The covariates sex and oxygen are not significant at level .05. This suggests that gender and oxygen do not play a significant role in determining the survival probability of a victim.

### APPENDIX: PROOFS

Before we present the proofs of the theorems, we first impose some regularity conditions. To this end, let us recall that  $q_j(s, y) = (\partial^j / \partial s^j)l\{g^{-1}(s), y\}$ . Note that  $q_k(s, y)$  is linear in  $y$  for fixed  $s$  such that

$$q_1[g\{m(u, \mathbf{x})\}, m(u, \mathbf{x})] = 0 \quad (A.1)$$

and

$$q_2[g\{m(u, \mathbf{x})\}, m(u, \mathbf{x})] = -\rho(u, \mathbf{x}), \quad (A.2)$$

where  $\rho(u, \mathbf{x})$  is defined in (5). Note that we use the same notation as in Section 2.

The regularity conditions are as follows:

C1. The function  $q_2(s, y) < 0$  for  $s \in \mathfrak{R}$  and  $y$  in the range of the response variable.

C2. The functions  $f_U(u), \Gamma(u), V(m(u, \mathbf{x})), V'(m(u, \mathbf{x}))$ , and  $g'''(m(u, \mathbf{x}))$  are continuous at the point  $u = u_0$ . Further, assume that  $f_U(u_0) > 0$  and  $\Gamma(u_0) > 0$ .

C3.  $K(\cdot)$  has a bounded support.

C4.  $a_j'(\cdot)$  is continuous in a neighborhood of  $u_0$  for  $j = 1, \dots, p$ .

C5.  $E\{\mathbf{X}|U = u\}$  is continuous at the point  $u = u_0$ .

C6.  $E(Y^4|U = u, \mathbf{X} = \mathbf{x})$  is bounded in a neighborhood of  $u = u_0$ .

Condition C1 guarantees that the local likelihood function (2) is concave. It is satisfied for the canonical exponential family with a canonical link. Note that condition C2 implies that  $q_1(\cdot, \cdot), q_2(\cdot, \cdot), q_3(\cdot, \cdot), \rho'(\cdot, \cdot)$ , and  $m'(\cdot, \cdot)$  are continuous.

### Proof of Theorem 1

Recall that  $\hat{\beta}_{\text{MLE}}$  maximizes (2). Let  $\bar{\eta}(u_0, u, \mathbf{x}) = \sum_{j=1}^p \{a_j(u_0) + a_j'(u_0)(u - u_0)\}x_j$  and

$$\beta^* = \gamma_n^{-1}(\beta_1 - a_1(u_0), \dots, \beta_p - a_p(u_0)),$$

$$h(\beta_{p+1} - a_1'(u_0), \dots, h(\beta_{2p} - a_p'(u_0)))^T,$$

where  $\gamma_n = (nh)^{-1/2}$ . It can be easily seen that  $\sum_{j=1}^p \{a_j + b_j(U_i - u_0)\}X_{ij} = \bar{\eta}(u_0, U_i, \mathbf{X}_i) + \gamma_n \beta^{*T} \mathbf{Z}_i$ , where  $\mathbf{Z}_i = (\mathbf{X}_i^T, ((U_i - u_0)/h)\mathbf{X}_i^T)^T$ . Then the local likelihood function  $l_n(\beta)$  defined in (2) becomes

$$l_n(\beta) = \frac{1}{n} \sum_{i=1}^n l[g^{-1}\{\bar{\eta}(u_0, U_i, \mathbf{X}_i) + \gamma_n \beta^{*T} \mathbf{Z}_i\}, Y_i] K_h(U_i - u_0),$$

which is a function of  $\beta^*$  denoted by  $l_n(\beta^*)$ . Let

$$\hat{\beta}^* = \gamma_n^{-1}(\hat{\beta}_1 - a_1(u_0), \dots, \hat{\beta}_p - a_p(u_0)),$$

$$h(\hat{\beta}_{p+1} - a_1'(u_0), \dots, h(\hat{\beta}_{2p} - a_p'(u_0)))^T.$$

Then  $\hat{\beta}^*$  maximizes  $l_n(\beta^*)$ , because  $\hat{\beta}$  maximizes (2). Equivalently,  $\hat{\beta}^*$  maximizes the normalized function

$$l_n^*(\beta^*) = \sum_{i=1}^n [l\{g^{-1}(\bar{\eta}_i(u_0) + \gamma_n \beta^{*T} \mathbf{Z}_i), Y_i\} - l\{g^{-1}(\bar{\eta}_i(u_0)), Y_i\}] K\{(U_i - u_0)/h\},$$

where  $\bar{\eta}_i(u_0) = \bar{\eta}(u_0, U_i, \mathbf{X}_i)$ .

We remark that condition C1 implies that  $l_n^*(\cdot)$  is concave in  $\beta^*$ . Using the Taylor expansion of  $l\{g^{-1}(\cdot), y\}$ , we have

$$l_n^*(\beta^*) = \mathbf{W}_n^T \beta^* + \frac{1}{2} \beta^{*T} \Delta_n \beta^* + \frac{\gamma_n^3}{6} \sum_{i=1}^n q_3\{\eta_i, Y_i\} (\beta^{*T} \mathbf{Z}_i)^3 K\{(U_i - u_0)/h\}, \quad (\text{A.3})$$

where

$$\mathbf{W}_n = \gamma_n \sum_{i=1}^n q_1\{\bar{\eta}_i(u_0), Y_i\} \mathbf{Z}_i K\{(U_i - u_0)/h\}, \quad (\text{A.4})$$

$$\Delta_n = \frac{\gamma_n^2}{2} \sum_{i=1}^n q_2\{\bar{\eta}_i(u_0), Y_i\} \mathbf{Z}_i \mathbf{Z}_i^T K\{(U_i - u_0)/h\},$$

and  $\eta_i$  is between  $\bar{\eta}_i(u_0)$  and  $\bar{\eta}_i(u_0) + \gamma_n \beta^{*T} \mathbf{Z}_i$ . Note that

$$(\Delta_n)_{ij} = (E\Delta_n)_{ij} + O_p\{\{\text{var}(\Delta_n)_{ij}\}^{1/2}\}.$$

Now the mean in the foregoing expression equals

$$E(\Delta_n) = h^{-1} E[q_2\{\bar{\eta}(u_0, U, \mathbf{X}), m(U, \mathbf{X})\} K\{(U - u_0)/h\} \mathbf{Z}\mathbf{Z}^T].$$

By a Taylor series expansion of  $\eta(u, \mathbf{x})$  with respect to  $u$  around  $|u - u_0| < h$  and (A.1), we have

$$\eta(u, \mathbf{x}) = \bar{\eta}(u_0, u, \mathbf{x}) + \frac{h^2(u - u_0)^2}{2} \eta_u''(u_0, \mathbf{x}) + o(h^2),$$

where  $\eta_u''(u, \mathbf{x}) = (\partial^2/\partial u^2)\eta(u, \mathbf{x}) = \sum_{j=1}^p a_j''(u)x_j$ , which implies that

$$q_1\{\bar{\eta}(u_0, u, \mathbf{x}), m(u, \mathbf{x})\} = \rho(u, \mathbf{x}) \frac{h^2(u - u_0)^2}{2} \eta_u''(u_0, \mathbf{x}) + o(h^2) \quad (\text{A.5})$$

and

$$q_2\{\bar{\eta}(u_0, u, \mathbf{x}), m(u, \mathbf{x})\} = -\rho(u, \mathbf{x}) + o(1). \quad (\text{A.6})$$

Then, using (A.2) and (A.6), we obtain

$$E(\Delta_n) \rightarrow -f_U(u_0) \begin{pmatrix} 1 & \mu_1 \\ \mu_1 & \mu_2 \end{pmatrix} \otimes \Gamma(u_0) = -\Delta, \quad (\text{A.7})$$

where  $\Gamma(u_0)$  is given in (4) and  $\Delta$  is defined in (7). Similar arguments show that  $\text{var}\{(\Delta_n)_{ij}\} = O\{(nh)^{-1}\}$ . Therefore,

$$\Delta_n = -\Delta + o_p(1). \quad (\text{A.8})$$

Because  $K(\cdot)$  is bounded,  $q_3(\cdot, \cdot)$  is linear in  $Y_1$  and  $E(|Y_1||U_1, \mathbf{X}_1) < \infty$ , the expected value of the absolute value of the last term in (A.3) is bounded by

$$O(n\gamma_n^3 E|q_3(\eta_1, Y_1)\mathbf{X}_1^3 K\{(U_1 - u_0)/h\}|) = O(\gamma_n) \quad (\text{A.9})$$

by condition C5. Therefore, the last term in (A.3) is of order  $O_p(\gamma_n)$ . This, in conjunction with (A.3), (A.7), and (A.8), implies that

$$l_n^*(\beta^*) = \mathbf{W}_n^T \beta^* - \frac{1}{2} \beta^{*T} \Delta \beta^* + o_p(1).$$

An application of the quadratic approximation lemma (see, e.g., Fan and Gijbels 1996, p. 210) leads to

$$\hat{\beta}^* = \Delta^{-1} \mathbf{W}_n + o_p(1), \quad (\text{A.10})$$

if  $\mathbf{W}_n$  is a sequence of stochastically bounded random vectors. The asymptotic normality of  $\hat{\beta}^*$  follows from that of  $\mathbf{W}_n$ . Hence it remains to establish the asymptotic normality of  $\mathbf{W}_n$ .

Note that the random vector  $\mathbf{W}_n$  is a sum of iid random vectors. To establish its asymptotic normality, it suffices to compute the mean and covariance matrix of  $\mathbf{W}_n$  and check the Lyapounov condition. To this end, by (A.5), we have

$$E(\mathbf{W}_n) = n\gamma_n E[q_1\{\bar{\eta}(u_0, U, \mathbf{X}), m(U, \mathbf{X})\} \mathbf{Z} K\{(U - u_0)/h\}] = \frac{h^2 f_U(u_0)}{2\gamma_n} \begin{pmatrix} \mu_2 \\ \mu_3 \end{pmatrix} \otimes \Gamma(u_0) \mathbf{a}''(u_0) \{1 + o(1)\}. \quad (\text{A.11})$$

Similarly, by (A.11) and the definition of  $q_1(\cdot, \cdot)$ , one has

$$\begin{aligned} \text{var}(\mathbf{W}_n) &= h^{-1} E[q_1^2\{\bar{\eta}(u_0, U, \mathbf{X}), Y\} \mathbf{Z}\mathbf{Z}^T K^2\{(U - u_0)/h\}] \\ &= f_U(u_0) \begin{pmatrix} \nu_0 & \nu_1 \\ \nu_1 & \nu_2 \end{pmatrix} \otimes \Gamma(u_0) \{1 + o(1)\} \\ &= \mathbf{\Lambda} + o(1), \end{aligned} \quad (\text{A.12})$$

where  $\mathbf{\Lambda}$  is defined in (8). By the Cramér–Wold device, to derive the asymptotic normality of  $\mathbf{W}_n$ , it suffices to show that for any unit vector  $\mathbf{d} \in \mathfrak{R}^{2p}$ ,

$$\{\mathbf{d}^T \text{var}(\mathbf{W}_n) \mathbf{d}\}^{-1/2} \{\mathbf{d}^T \mathbf{W}_n - \mathbf{d}^T E(\mathbf{W}_n)\} \xrightarrow{D} N(0, 1). \quad (\text{A.13})$$



This, in conjunction with (A.10), (A.11), and (A.12), implies that

$$\hat{\beta}^* - \frac{(nh^5)^{1/2}}{2} \Delta^{-1} f_U(u_0) \begin{pmatrix} \mu_2 \\ \mu_3 \end{pmatrix} \otimes \Gamma(u_0) \mathbf{a}''(u_0) \{1 + o(1)\} \xrightarrow{D} N(\mathbf{0}, \Delta^{-1} \Lambda \Delta^{-1}). \quad (\text{A.14})$$

Therefore, the assertion in (6) holds true. To prove (A.12), we need only check Lyapounov's condition for that sequence. To do so, let  $\xi_i = q_1 \{ \bar{\eta}_i(u_0), Y_i \} \mathbf{d}^T \mathbf{Z}_i K \{ (U_i - u_0)/h \}$ . Then  $\mathbf{d}^T \mathbf{W}_n = \gamma_n \sum_{i=1}^n \xi_i$ . It suffices to show that  $n\gamma_n^3 E|\xi_1|^3 \rightarrow 0$  as  $n \rightarrow \infty$ . Similar to (A.8), one can show that  $n\gamma_n^3 E|\xi_1|^3 = O(\gamma) \rightarrow 0$ . If  $K(\cdot)$  is symmetric, then  $\mu_1 = 0$ , so that (9) holds true. This completes the proof of the theorem.

**Proof of Theorem 2**

Recall that

$$l_n(\beta) = \frac{1}{n} \sum_{i=1}^n l \left\{ g^{-1} \left( \sum_{j=1}^p (a_j + b_j(U_i - u_0)) X_{ij} \right), Y_i \right\} \times K_h(U_i - u_0).$$

For any  $\tilde{\beta}$  satisfying  $\mathbf{H}(\tilde{\beta} - \beta) = O_p(h^2 + (nh)^{-1/2})$ , one can easily show that

$$\begin{aligned} \mathbf{H}^{-1} l_n''(\tilde{\beta}) \mathbf{H}^{-1} &= \mathbf{H}^{-1} l_n''(\beta) \mathbf{H}^{-1} + o_p(1) \\ &= \frac{1}{n} \sum_{i=1}^n q_2 \{ \tilde{\mathbf{Z}}_i^T \beta, Y_i \} \mathbf{H}^{-1} \tilde{\mathbf{Z}}_i \tilde{\mathbf{Z}}_i^T \mathbf{H}^{-1} K_h(U_i - u_0) \\ &\quad + o_p(1), \end{aligned} \quad (\text{A.15})$$

where  $\tilde{\mathbf{Z}}_i = (\mathbf{X}_i^T, (U_i - u_0)\mathbf{X}_i^T)^T$ . By computing the mean and variance of  $\mathbf{H}^{-1} l_n''(\tilde{\beta}) \mathbf{H}^{-1}$ , we obtain

$$\begin{aligned} \mathbf{H}^{-1} l_n''(\tilde{\beta}) \mathbf{H}^{-1} &= E \left[ q_2 \{ \tilde{\mathbf{Z}}^T \beta, Y \} \begin{pmatrix} 1 \\ (U - u_0)/h \end{pmatrix} \otimes \mathbf{X}\mathbf{X}^T K_h(U - u_0) \right] + o_p(1) \\ &= E \left[ q_2 \{ \tilde{\mathbf{Z}}^T \beta, m(U, \mathbf{X}) \} \begin{pmatrix} 1 \\ (U - u_0)/h \end{pmatrix} \otimes \mathbf{X}\mathbf{X}^T K_h(U - u_0) \right] + o_p(1) \\ &= -\Delta + o_p(1), \end{aligned} \quad (\text{A.16})$$

where  $\Delta$  is defined in (7). Recall that  $\hat{\beta}_{OS} = \hat{\beta}_0 - \{ l_n''(\hat{\beta}_0) \}^{-1} l_n'(\hat{\beta}_0)$ ; see (3). By the Taylor expansion, we have

$$l_n'(\hat{\beta}_0) = l_n'(\beta) + l_n''(\tilde{\beta}^*)(\hat{\beta}_0 - \beta),$$

where  $\tilde{\beta}^*$  lies between  $\beta$  and  $\hat{\beta}_0$  and hence satisfies  $\mathbf{H}(\tilde{\beta}^* - \beta) = O_p(h^2 + (nh)^{-1/2})$ . Then, some algebraic computations show that

$$\begin{aligned} \mathbf{H}(\hat{\beta}_{OS} - \beta) &= \mathbf{H}(\hat{\beta}_0 - \beta) - \mathbf{H} \{ l_n''(\hat{\beta}_0) \}^{-1} \mathbf{H} \mathbf{H}^{-1} l_n'(\hat{\beta}_0) \\ &= [\mathbf{I} - \mathbf{H} \{ l_n''(\hat{\beta}_0) \}^{-1} \mathbf{H} \mathbf{H}^{-1} l_n''(\tilde{\beta}^*) \mathbf{H}^{-1}] \mathbf{H}(\hat{\beta}_0 - \beta) \\ &\quad - \mathbf{H} \{ l_n''(\hat{\beta}_0) \}^{-1} \mathbf{H} \mathbf{H}^{-1} l_n'(\beta). \end{aligned} \quad (\text{A.17})$$

Therefore, by (A.16) and (A.17), we have

$$\mathbf{H}(\hat{\beta}_{OS} - \beta) = \Delta^{-1} \mathbf{H}^{-1} l_n'(\beta) \{1 + o_p(1)\} + o_p(h^2 + (nh)^{-1/2}),$$

which, in conjunction with (A.4), (A.10), (A.13), and (A.14), implies that

$$\sqrt{nh} \mathbf{H}(\hat{\beta}_{OS} - \beta) = \Delta^{-1} \mathbf{W}_n + o_p(1) = \hat{\beta}^* + o_p(1). \quad (\text{A.18})$$

Therefore,  $\hat{\beta}_{OS}$  has the same asymptotic distribution as  $\hat{\beta}_{MLE}$ .

[Received March 1999. Revised December 1999.]

**REFERENCES**

Bickel, P. J. (1975), "One-Step Huber Estimates in Linear Models," *Journal of the American Statistical Association*, 70, 428-433.

Brumback, B., and Rice, J. (1998), "Smoothing Spline Models for the Analysis of Nested and Crossed Samples of Curves," *Journal of the American Statistical Association*, 93, 961-976.

Cai, Z., Fan, J., and Yao, Q. (1998), "Functional-Coefficient Regression Models for Nonlinear Time Series," *Journal of the American Statistical Association*, to appear in December 2000.

Cai, Z., and Tiwari, R. C. (in press), "Application of a Local Linear Autoregressive Model to BOD Time Series," *Environmetrics*.

Carroll, R. J., Fan, J., Gijbels, I., and Wand, M. P. (1997), "Generalized Partially Linear Single-Index Models," *Journal of the American Statistical Association*, 92, 477-489.

Carroll, R. J., Ruppert, D., and Welsh, A. H. (1998), "Local Estimating Equations," *Journal of the American Statistical Association*, 93, 214-227.

Chen, H. (1988), "Convergence Rates for Parametric Components in a Partly Linear Model," *The Annals of Statistics*, 16, 136-146.

Chen, R., and Tsay, R. S. (1993), "Functional-Coefficient Autoregressive Models," *Journal of the American Statistical Association*, 88, 298-308.

Cleveland, W. S., Grosse, E., and Shyu, W. M. (1992), "Local Regression Models," in *Statistical Models in S*, (eds. J. M. Chambers and T. J. Hastie, Pacific Grove, CA: Wadsworth & Brooks/Cole, pp. 309-376.

Fan, J. (1993), "Local Linear Regression Smoothers and Their Minimax," *The Annals of Statistics*, 21, 196-216.

Fan, J., and Chen, J. (1999), "One-Step Local Quasi-Likelihood Estimation," *Journal of the Royal Statistics Society, Ser. B*, 61, 927-943.

Fan, J., and Gijbels, I. (1996), *Local Polynomial Modeling and Its Applications*, London: Chapman and Hall.

Fan, J., Zhang, C., and Zhang, J. (1999), "Sieve Likelihood Ratio Statistics and Wilks Phenomenon," technical report, UCLA, Dept. of Statistics.

Fan, J., and Zhang, J. (2000), "Functional Linear Models for Longitudinal Data," *Journal of the Royal Statistical Society, Ser. B*, 62, 303-322.

Fan, J., and Zhang, W. (1999), "Statistical Estimation in Varying-Coefficient Models," *The Annals of Statistics*, 27, 1491-1518.

Green, P. J., and Silverman, B. W. (1994), *Nonparametric Regression and Generalized Linear Models: A Robust Penalty Approach*, London: Chapman and Hall.

Hastie, T. J., and Tibshirani, R. J. (1993), "Varying-Coefficient Models" (with discussion), *Journal of the Royal Statistical Society, Ser. B*, 55, 757-796.

Hong, Y., and Lee, T.-H. (1999), "Inference and Forecast of Exchange Rates via Generalized Spectrum and Nonlinear Times Series Models," unpublished manuscript.

Hoover, D. R., Rice, J. A., Wu, C. O., and Yang, L. P. (1998), "Nonparametric Smoothing Estimates of Time-Varying Coefficient Models With Longitudinal Data," *Biometrika*, 85, 809-822.

Kauermann, G., and Tutz, G. (1999), "On Model Diagnostics Using Varying-Coefficient Models," *Biometrika*, 86, 119-128.

Lee, T.-H., and Ullah, A. (1999), "Nonparametric Bootstrap Tests for Neglected Nonlinearity in Time Series Regression Models," unpublished manuscript.

Lehmann, E. L. (1983), *Theory of Point Estimation*, Pacific Grove, CA: Wadsworth & Brooks/Cole.

McCullagh, P., and Nelder, J. A. (1989), *Generalized Linear Models* (2nd ed.), London: Chapman and Hall.

Ruppert, D., and Wand, M. P. (1994), "Multivariate Weighted Least Squares Regression," *The Annals of Statistics*, 22, 1346-1370.

Seifert, B., and Gasser, T. (1996), "Finite-Sample Variance of Local Polynomial: Analysis and Solutions," *Journal of the American Statistical Association*, 91, 267-275.

Speckman, P. (1988), "Kernel Smoothing in Partial Linear Models," *Journal of the Royal Statistical Society, Ser. B*, 50, 413-436.